

Ancient DNA from an extinct native duck reveals how far birds flew to make New Zealand home

May 20 2024, by Nic Rawlence and Alexander Verry



Auckland Island Merganser, *Mergus australis*, collected June 1902, Auckland Islands, New Zealand. Credit: [Te Papa \(OR.001357\)](#), [CC BY](#)

Ask a bird lover if they have heard of the extinct [giant moa](#) or its ancient

predator, [Haast's eagle](#), and the answer will likely be yes. The same can't be said of New Zealand's extinct, but equally unique, [mergansers](#)—a group of fish-eating ducks with a serrated bill.

The only southern hemisphere representatives of this group are the critically endangered [Brazilian merganser](#) and those from the [New Zealand region](#), which are now extinct.

Unlike some of New Zealand's other [extinct birds](#), the biological heritage of our enigmatic mergansers is shrouded in mystery. But our [new research](#) on the extinct Auckland Island merganser is changing the way we think about the origins of New Zealand's birds. Did the ancestors of the merganser come from South America or the northern hemisphere—and when did they arrive?

Lost to humans and pests

[Mergansers were spread across](#) the three main islands of New Zealand at the time of Polynesian arrival in the 13th century, as well as the Auckland Islands to the south and the [Chatham Islands](#) to the east.

Over-hunting, [habitat destruction](#), and predation from the Pacific rat and [Polynesian dog](#) resulted in the extinction of mergansers on the New Zealand mainland and the Chatham Islands. By the time Europeans arrived in the 17th century, mergansers were restricted to an isolated population on the sub-Antarctic Auckland Islands.

European discovery of the Auckland Islands in 1806 led to a formal description of the [Auckland Island merganser](#) in 1841. However, European discovery brought new predators like pigs and cats.

Mergansers were also sought after as specimens for the museum trade. The last known Auckland Island merganser was shot and collected in

1902, a mere 61 years after its discovery.

Only recently have merganser bones from the Chatham Islands been described as a distinct species. The distinction of the [Chatham Island merganser](#) hints at greater merganser diversity in the New Zealand region than previously thought. It is possible the merganser bones found on mainland New Zealand may be another species, but only ongoing research will be able to confirm this.

From the northern hemisphere to NZ

The extinction of mergansers from the New Zealand region has meant their evolutionary history has remained a mystery. Did their ancestors, and those of the Brazilian merganser, arrive via independent colonization events from the northern hemisphere? Or was there a single push into the southern hemisphere, followed by subsequent divergence events?

To find out more, we [sequenced ancient DNA](#) from an Auckland Islands merganser and a Brazilian merganser. This allowed us to reconstruct the evolutionary history of the wider group.

We found mergansers originated in the northern hemisphere, diverging from their closest relatives some 18 million years ago, before rapidly evolving into several different species between 14 and seven million years ago.

The mergansers from the New Zealand region are most closely related to the northern hemisphere [common merganser](#). Their ancestors arrived here at least seven million years ago in a separate colonization event to the one that gave rise to the Brazilian merganser.

Further [genetic research](#) is currently underway. The goal is to reconstruct the [evolutionary history](#) of mergansers within the New Zealand region.

The global origins of New Zealand's birds

Many New Zealanders believe the country's [native birds originate](#) from Australia. Increasingly though, genetic and paleontological research shows a number of our feathered friends hail from further afield.

Kiwi are most closely related to the extinct [elephant birds of Madagascar](#), for example. And the extinct [adzebill](#) is related to [flufftails, also from Madagascar](#). The extinct moa is most closely related to the [tinamou from South America](#).

The long journey of [blue-eyed shags](#) started in South America, with the birds island hopping via Antarctica and the sub-Antarctic islands to New Zealand. Mergansers arriving from the northern hemisphere add another piece to the puzzle.

It is possible that fossils of extinct mergansers (and other birds with distant geographic origins) will be discovered as paleontologists increasingly focus on previously neglected and newly discovered southern hemisphere fossil deposits.

Only then, combined with the power of ancient DNA, will we be able to fully understand how New Zealand's dynamic geological, climatic and human history has influenced the colonization and diversification of birds on this isolated South Pacific archipelago.

This article is republished from [The Conversation](#) under a Creative Commons license. Read the [original article](#).

Provided by The Conversation

Citation: Ancient DNA from an extinct native duck reveals how far birds flew to make New Zealand home (2024, May 20) retrieved 12 September 2024 from <https://phys.org/news/2024-05-ancient-dna-extinct-native-duck.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--