

# Investigating coal emissions reductions and mortality in China

May 1 2024




Figure 1. Annual PM<sub>2.5</sub> concentrations in BTH and surrounding areas from 2013 to 2018

Annual PM<sub>2.5</sub> pollution in the Beijing-Tianjin-Hebei area from 2013 to 2018, showing a decreasing trend by year, especially in the southern area of Hebei Province. Credit: Li et al

In 2012, China was the largest consumer of coal in the world. In 2013, the State Council of China issued the "China National Action Plan on Air Pollution Prevention and Control," aiming to reduce emissions from coal combustion. The plan included renovations to small residential coal heating stoves as well as retrofits for coal-fired power plants, among other initiatives.

Annual coal consumption fell between 2013 and 2017, which led to observed dramatic decreases in mean daily fine particulate matter (PM<sub>2.5</sub>) levels.

In 2018, a new program, known as "Three-Year Action Plan for Winning the Blue Sky Defense Battle," began, and in the same year, PM<sub>2.5</sub> concentrations were further reduced by 9.3% from 2017 levels.

In this context, Xiaoming Shi and colleagues used accountability analysis to assess whether the acute health effects of PM<sub>2.5</sub> changed from 2013 to 2018 in the Beijing-Tianjin-Hebei area, which was the most heavily polluted region. The study is [published](#) in the journal *PNAS Nexus*.




Figure 3. The effect of PM<sub>2.5</sub> on mortality in BTH and surrounding areas (A) and control areas (B) during 2013-2015 and 2016-2018, and exposure-response curves (curves and shading indicate RR and 95% CI) in different study periods (C)

\*Indicates that the difference between the groups was statistically significant

The percentage changes in mortality for total, circulatory, and respiratory diseases associated with each 10 µg/m<sup>3</sup> increase in PM<sub>2.5</sub> in the Beijing-Tianjin-Hebei area. Credit: Li et al

The acute effects of PM<sub>2.5</sub> were significantly decreased for total and circulatory mortality. A 10 µg/m<sup>3</sup> increase in PM<sub>2.5</sub> concentrations was associated with an increase of 0.16% and 0.02% in mortality from 2013 to 2015 and from 2016 to 2018, respectively.

According to the authors, the reduced exposure-response relationship could be due to changes in particle toxicity, as PM<sub>2.5</sub> from other sources may be less toxic than PM<sub>2.5</sub> from coal. Behavioral changes to avoid air pollution may also have influenced the outcomes, the authors add.

**More information:** Tiantian Li et al, Accountability analysis of health benefits related to National Action Plan on Air Pollution Prevention and Control in China, *PNAS Nexus* (2024). [DOI: 10.1093/pnasnexus/pgae142](https://doi.org/10.1093/pnasnexus/pgae142)

Provided by PNAS Nexus

Citation: Investigating coal emissions reductions and mortality in China (2024, May 1) retrieved 16 May 2024 from <https://phys.org/news/2024-05-coal-emissions-reductions-mortality-china.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.