

New feather mite species discovered on the endangered Okinawa rail

May 17 2024



Left: male. Right: female. Credit: Dr. Tsukasa Waki



A research group led by Dr. Tsukasa Waki of Toho University and Professor Satoshi Shimanono of Hosei University have discovered a new mite species, Metanalges agachi, which is thought to clean the feathers of the endangered Okinawa rail, endemic to the northern part of Okinawa Island, Japan.

It has been 43 years since the endangered Okinawa rail was first described as a <u>new species</u>. The scientific name "agachi," derived from the affectionate local name "Agachi," meaning "impatient or hasty person," which the Okinawa rail has been called in this region for generations, was chosen for this species. Assigning this scientific name to the <u>mites</u> found on the Okinawa rail has ensured that the affectionate nickname of the Okinawa rail in this region has been preserved for eternity.

This research was <u>published</u> online in the journal *Species Diversity* on April 24 2024.

The Okinawa rail is an endemic species found in the northern part of Okinawa Island and has been designated a national natural monument. It is classified as "Critically Endangered" in the Ministry of the Environment's Red List and "Endangered" in the IUCN Red List. The Okinawa rail has been threatened with extinction due to predation by <u>invasive species</u>, such as mongooses, because of its flightless nature. Owing to various conservation efforts, <u>population numbers</u> have recently shown signs of recovery.





This bird is the only host of Meatanalges agachi. Credit: Yambaru Ranger Office

The feather mite, Metanalges agachi, is found on the feathers of the Okinawa rail and feeds on the oil, fungi, bacteria, and other debris that accumulate on it. It is considered a mutualistic symbiont that helps clean the bird's feathers, as it does not feed on living bird tissue or blood.

The research group examined more than 300 frozen specimens of the Okinawa rail collected and stored between 2008 and 2020. This led to the discovery of a new mite species, Metanalges agachi, found only on



the Okinawa rail. Without the Okinawa rail, this mite species will not survive. Therefore, like their hosts, these mites are also at risk of <u>extinction</u>, potentially even more so.

More information: Tsukasa Waki et al, A New Feather Mite Species of the Genus Metanalges (Acariformes: Analgidae) from the Okinawa Rail, Hypotaenidia okinawae (Gruiformes: Rallidae), in Okinawa Island, Japan, *Species Diversity* (2024). <u>DOI: 10.12782/specdiv.29.161</u>

Provided by Toho University

Citation: New feather mite species discovered on the endangered Okinawa rail (2024, May 17) retrieved 23 June 2024 from <u>https://phys.org/news/2024-05-feather-mite-species-endangered-okinawa.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.