


Genetic mutation responsible for new coat pattern in cats in Finland identified

May 23 2024, by Bob Yirka



Salmiak coloring in cats. Prominent features of the coloring are: "tuxedo" (a.k.a. bicolor) white spotting in the absence of white spotting alleles (W_s , g), and additional gradation of the pigment within hairs of primary color toward no pigmentation at the tips in the body, legs and tail. Additionally, there is primary colored spotting in the white areas of the front legs and chest, more intense

coloring in the scapular region, and a very pale tip of the tail. Photo credits: (a) Ari Kankainen and (b–e) courtesy of the cat owners. Credit: *Animal Genetics* (2024). DOI: 10.1111/age.13438

A team of geneticists and animal welfare specialists from the U.K. and one from the U.S. has identified the genetic mutation responsible for a newly discovered cat coat color pattern in Finland. In their [study](#), published in the journal *Animal Genetics*, the group sequenced the genomes of two of the cats.

In 2007, people in Finland began to notice that some [feral cats](#) had a distinctively colored coat. Instead of what would be a typical tuxedo pattern, the cats had hairs that were black at the base and grew progressively whiter toward the tip. The pattern soon became known as the "Finnish mutation" as some were adopted and brought into homes. In this new study, the research team explored the genetic mutation that was responsible for the change.

The researchers gave the new coat pattern an official name—they dubbed it salmiak, after a popular Finnish licorice. They next put word out that they were looking for cats with the new coat pattern, asking volunteers to bring them in to be tested.

Initial testing of the cats turned up negative for mutations that are known to impact white fur coloration. That led the team to conduct whole genome sequencing for two of the cats. They found a mutation at a site near the KIT gene that had previously been associated with several white hair patterns in housecats—they also noted that a span of the sequence was missing downstream.

Next, the research group verified they had found the [variant](#) they were looking for—they looked for it in 181 cats and found it in three of them that had the salmiak coat. All three had also inherited it from both of their parents. They also found three other cats that had the variant but not the salmiak coat—they had inherited the mutation from just one parent.

The research team suggests they have identified the recessive mutation responsible for the development of the new, unique coat pattern.

More information: Heidi Anderson et al, A new Finnish flavor of feline coat coloration, "salmiak," is associated with a 95-kb deletion downstream of the KIT gene, *Animal Genetics* (2024). [DOI: 10.1111/age.13438](#)

© 2024 Science X Network

Citation: Genetic mutation responsible for new coat pattern in cats in Finland identified (2024, May 23) retrieved 25 June 2024 from <https://phys.org/news/2024-05-genetic-mutation-responsible-coat-pattern.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--