

## **Researchers shed light on how key ingredient** for life may form in space

May 16 2024



This graphic depicts methanol's chemical structure (CH<sub>3</sub>OH) breaking down into hydroxymethylene (HCOH), a crucial precursor to the building blocks of life. Credit: Leah Dodson and Emily Hockey.

A team led by University of Maryland chemists discovered a new way to create carbenes, a class of highly reactive yet notoriously short-lived and unstable molecules. Involved in many high-energy chemical reactions such as the creation of carbohydrates, carbenes are crucial precursors to the building blocks of life on Earth—and possibly in space.

The scientists successfully formed a carbene called hydroxymethylene



(HCOH) by breaking down methanol (a common alcohol found in many industrial chemicals like formaldehyde) with pulses of ultraviolet radiation. The results were published in <u>a paper</u> on May 14, 2024, in the *Journal of the American Chemical Society*.

"It's surprising to see this carbene come from such a commonplace molecule like methanol—we have squirt bottles of it in labs everywhere," said Leah Dodson, an assistant professor of Chemistry and Biochemistry at UMD and senior author of the paper.

"193-nanometer wavelength UV lasers are also fairly standard. This means that carbenes could be naturally forming in places like space, where there is a lot of methanol and ultraviolet radiation. And further reactions of carbenes formed in space through this process could lead to biomolecules that make up life."

The paper's findings reveal clues about the mechanisms behind carbene formation and reaction on Earth, leading to a better understanding of the molecule's potential to create sugars necessary for life.

"There's established research that suggests that HCOH can react to form simple sugars, including some that have previously been detected in space," said the study's lead author Emily Hockey. "We think it's possible that this carbene, since it comes from a molecule that's so ubiquitous in space and can be detected anywhere, is the missing piece bridging gaps in our knowledge of how methanol and simple sugars can lead to bigger, more advanced biomolecules."





Dodson (left) and Hockey (right) observing data at the Advanced Light Source research facility at Lawrence Berkeley National Laboratory in Berkeley, California, where they conducted their experiments with methanol and ultraviolet radiation. Credit: Leah Dodson and Emily Hockey.

Due to their super-reactivity, carbene molecules usually have very short lifetimes. These characteristics make carbenes generally difficult for scientists to generate and observe, which has limited deep understanding of the molecule. But the UMD team's novel method of producing carbenes allowed them to study the molecules closely enough to see their formation and decay over millisecond timescales. The researchers were surprised to discover that HCOH reacted relatively slowly with oxygen at room temperature.



"When we looked at HCOH's reactivity in our room temperature system, we saw that it decayed within 15 milliseconds," Hockey explained. "What's interesting is that because carbenes are thought to be a super reactive species, it's reasonable to assume that this carbene would react so quickly to something like oxygen that it's impossible to catch. But that's not what happened. Although the <u>carbene</u> was decaying faster and faster when exposed to oxygen, it was slow enough that we were still able to observe that decay."

The researchers believe that their method of producing and studying carbenes will help astronomers and astrochemists gain new insights into the origins of life and how life in space may have evolved differently from life on Earth. They hope to build on their findings by looking closer at what happens during methanol's breakdown and quantifying the different products yielded by methanol's reaction to UV light.

"We know that carbenes like HCOH are formed during our process, but we'd like to dig deeper into what percentage of it ends up as formaldehyde, methylene or other hydrocarbon radicals, for example," Hockey explained. "We originally thought all the products would be methoxy radicals but our experiments show that the process and the resulting products are more complicated than our original assumptions."

Knowing the types and amount of products created by breaking down methanol with UV radiation would provide astronomers and astrochemists with a more accurate outlook on astrophysical objects and how they evolved over billions of years.

"If the existing data on what is produced from <u>methanol</u> photodissociation are wrong, then the models being propagated will be incorrect as well—and our understanding of how life evolved from these molecules could also be compromised," Dodson said. "Our follow-up work will hopefully lay the groundwork for those types of simulations."



**More information:** Emily K. Hockey et al, Direct Observation of Gas-Phase Hydroxymethylene: Photoionization and Kinetics Resulting from Methanol Photodissociation, *Journal of the American Chemical Society* (2024). DOI: 10.1021/jacs.4c03090

## Provided by University of Maryland

Citation: Researchers shed light on how key ingredient for life may form in space (2024, May 16) retrieved 26 June 2024 from <u>https://phys.org/news/2024-05-key-ingredient-life-space.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.