


# Partial skeleton of a previously unknown medium-sized theropod dinosaur found in Siberia

May 28 2024, by Bob Yirka



*Kiyacursor longipes* gen. et sp. nov., referred specimen PIN 329/16 (b) and holotype KOKM 5542 (c–j). Shestakovo 1, Kemerovo oblast—Kuzbass, Russia; Ilek Formation, Lower Cretaceous (Aptian). Composite reconstruction including the known elements (by A. A. Atuchin). Credit: *Proceedings of the Royal Society B: Biological Sciences* (2024). DOI: 10.1098/rspb.2024.0537

Study of a partial skeleton found embedded in a rock has resulted in the


discovery of a new species of dinosaur. Using a variety of technology and techniques, researchers affiliated with several institutions in the Russian Federation found that the fossil once belonged to a previously unknown dinosaur they have named *Kiyacursor longipes*.

The team has published the [findings](#) in the journal *Proceedings of the Royal Society B: Biological Sciences*.

The rock was found to have recently fallen down a rocky cliffside in a part of western Siberia along the Kiya River due to natural erosion. The [fossilized bones](#) were sticking out of it, revealing its [ancient history](#).

The researchers studied the fossils visually and through use of X-ray and computed tomography. The research team also collected samples from some of the fossils and studied them using a microscope. As part of their analysis, they found that the fossilized skeleton was approximately 113 to 121 million years old. They identified ribs, vertebrae, part of a shoulder, both feet and both legs.

The dinosaur was a noasaurid ceratosaur, a group of bipedal, non-avian swift runners with two small legs that would have been tucked up at the top. The finding, they note, extends the range of Ceratosauria in Asia by approximately 40 million years.



*Kiyacursor longipes* gen. et sp. nov., KOKM 5542, fragmentary postcranial skeleton, holotype. Shestakovo 1, Kemerovo oblast—Kuzbass, Russia; Ilel Formation, Lower Cretaceous (Aptian). (a) Segmented surface visualization based on X-ray computed tomography scans. (b) Screenshot of photogrammetric model. Credit: *Proceedings of the Royal Society B: Biological Sciences* (2024). DOI: 10.1098/rspb.2024.0537

In measuring the length of the bones, the researchers found the dinosaur had what they describe as "unique hind-limb proportions" as compared to its known relatives, a feature that would have given the dinosaur better cursorial ability. It also had what they describe as "ostrich-like" feet, with its third toe extended—a feature not seen in any of its relatives.

They estimate that their *K. longipes* specimen was approximately 2.5

meters long (from nose to tail) when alive. They also found evidence that the medullary cavity inside the fossils that had once held bone marrow had ceased expanding, suggesting it was a mature adult, though not likely more than 3 years old at the time of its death.

**More information:** Alexander O. Averianov et al, The last ceratosaur of Asia: a new noosaurid from the Early Cretaceous Great Siberian Refugium, *Proceedings of the Royal Society B: Biological Sciences* (2024). [DOI: 10.1098/rspb.2024.0537](https://doi.org/10.1098/rspb.2024.0537)

© 2024 Science X Network

Citation: Partial skeleton of a previously unknown medium-sized theropod dinosaur found in Siberia (2024, May 28) retrieved 25 June 2024 from <https://phys.org/news/2024-05-partial-skeleton-previously-unknown-medium.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.