

Study explains regional differences in bird diversity in agroforestry systems

May 21 2024



Study areas in two contrasting landscapes with different dominant forest types in Peru. Credit: *Conservation Science and Practice* (2024). DOI: 10.1111/csp2.13123

The diversity and ecological functionality of bird communities in tropical agroforestry systems are shaped by the surrounding landscape, in



particular the extent and composition of the forest. An international research team led by the University of Göttingen has now investigated the composition and ecological traits of bird communities in 23 cocoa agroforestry systems in Peru.

The Universities of Würzburg and Vienna and the Alliance Biodiversity International in Peru were also involved in the study. The scientists found very different results depending on the region, and therefore emphasize the importance of tailoring <u>agroforestry</u> management strategies according to the region. The results are <u>published</u> in the journal *Conservation Science and Practice*.

The <u>cocoa</u> agroforestry systems studied are located in two contrasting Peruvian regions: one with seasonally dry tropical forests and one with subtropical humid rainforests along the Andes.

In both regions, cocoa agroforestry systems exist at different distances from natural forest, and increasing distance reduces the biodiversity in these agroforestry systems. The two regions differ greatly in their climate, as well as the structure and complexity of the forest vegetation.

"We found a much higher richness of bird species in the humid subtropical forest than in the dry forest—179 species compared to just 64," says first author Dr. Carolina Ocampo-Ariza from the University of Göttingen.

"In the dry forest landscapes, distance from the natural forest strongly influences the composition of bird communities, but less so the proportion of creatures that eat insects, which is a particularly important factor for biological pest control in cocoa farms."

Compared with cocoa farms that were right next to natural forest, if there was a distance of one kilometer, farms in the subtropical forest



experienced a decrease of 27% in the proportion of insects in the birds' diet; whereas for farms in the dry forest, the decrease was just 3%.

"These large regional differences in Peru show that agroforestry management needs to be adapted regionally to support optimal preservation and promotion of bird diversity and its ecosystem services. This is particularly true for biological pest control, which is important for cocoa yields," says Ocampo-Ariza.

"The proximity of cocoa cultivation to the nearest <u>natural forest</u> plays a special role for bird diversity and its ecosystem services."

Co-author Professor Teja Tscharntke adds, "Forested landscapes are crucial for the biodiversity in agroforestry systems. In addition, it is necessary to maintain and encourage a large number of older shadeproviding trees, in addition to complex vegetation, in the farming systems."

More information: Carolina Ocampo-Ariza et al, Regional differences of functional and taxonomic bird diversity in tropical agroforests of Peru, *Conservation Science and Practice* (2024). DOI: 10.1111/csp2.13123

Provided by University of Göttingen

Citation: Study explains regional differences in bird diversity in agroforestry systems (2024, May 21) retrieved 20 June 2024 from <u>https://phys.org/news/2024-05-regional-differences-bird-diversity-agroforestry.html</u>



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.