


# **Unlocking the secrets of supercritical fluids: Study offers insights into a hybrid state of matter**

May 22 2024

---



Supercritical fluids are crucial to understand giant planets such as Jupiter and Neptune, where similar states of matter may reign. Credit: ILL

A study now [published](#) in *Nature Communications* brings remarkable insights into the enigmatic behavior of supercritical fluids, a hybrid state of matter occupying a unique space between liquids and gases, and

arising in domains that go from the pharmaceutical industry to planetary science. The obtained results are at the limit of current experimental possibilities and could only be obtained in a high flux neutron source such as the Institut Laue-Langevin (ILL).

A liquid or gaseous substance pushed beyond its critical point (i.e., beyond the temperature and pressure at which the distinction between liquid and gas can no longer be made) is called a supercritical fluid. Still little-known and defying conventional classifications, [supercritical fluids](#) possess the ability to effuse like a gas while dissolving materials like a liquid.

This duality has made them invaluable in a myriad of industrial applications, from pharmaceutical processing to decaffeinating coffee beans. On the other hand, they are crucial to understand [giant planets](#) such as Jupiter and Neptune, where similar states of matter may reign.

An international team of researchers from Sapienza University (Rome, Italy) ILL (Grenoble, France), Ecole Polytechnique Federal (Lausanne, Switzerland), CNRS (France) and CNR (Italy) obtained [experimental proof](#) that [molecular diffusion](#) in a superfluid switches from gaseous-like behavior to liquid-like behavior across the so-called Widom line (a thermodynamic line that extends the saturated vapor curve above the [critical point](#)). The transition is gradual within a narrow pressure range.

The team investigated the diffusion of molecules within a supercritical fluid—a crucial parameter reflecting the mobility of molecules within the fluid—with a fundamental question in mind: can we pinpoint a region of pressure-temperature where the behavior of a supercritical fluid goes from gas-like to liquid-like? While [theoretical models](#) have proposed various such transition boundaries (among them the Widom

line) experimental validation had remained, until now, elusive.

This result was obtained through challenging, high-pressure, quasi-elastic neutron scattering (QENS) experiments on supercritical methane conducted at the ILL, in Grenoble. At the ILL, neutrons are used to explore materials and processes in all possible ways in a very wide range of domains.

In this study, a neutron beam was sent onto a cell containing methane in supercritical conditions. The intensity of the neutron beam scattered by the sample was measured as a function of the energy exchanged in the range of interest (i.e., in the energy range where molecular diffusion phenomena within matter occur, the so-called quasi-elastic regime).

The measurements took place at constant temperature  $T=200$  K (above the critical  $T=190$  K) varying the pressure of the methane from a few bars up to very high pressures (reaching nearly 3 Kbar; the critical pressure is  $P=45$  bar). The experiments were conducted on ILL instrument IN6-SHARP.

The authors underline the strikingly clear experimental evidence, "While at pressures lower than approximately 50 bar the signal of the diffusion dynamics typical of gaseous systems is observed, we have been able to observe that as the [pressure](#) increases above that, the signal evolves progressively until it takes on the typical shape of liquids," explains author Alessio De Francesco (researcher at CNR and ILL).

The result was made possible thanks to the high flux [neutron](#) source and the unique experimental support facilities available available at the ILL. "These measurements are at the limits of current experimental possibilities, and were unthinkable until a few years ago," adds Ferdinando Formisano (researcher at CNR and ILL).

"As often happens in research, having opened a door means seeing new avenues to explore, and this objective can only be pursued thanks to access to large research facilities."

**More information:** Umbertoluca Ranieri et al, Crossover from gas-like to liquid-like molecular diffusion in a simple supercritical fluid, *Nature Communications* (2024). DOI: [10.1038/s41467-024-47961-7](https://doi.org/10.1038/s41467-024-47961-7)

Provided by Institut Laue-Langevin

Citation: Unlocking the secrets of supercritical fluids: Study offers insights into a hybrid state of matter (2024, May 22) retrieved 16 June 2024 from <https://phys.org/news/2024-05-secrets-supercritical-fluids-insights-hybrid.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.