

# Summers warm up faster than winters, fossil shells from Antwerp show

May 15 2024

---



A fossil specimen of *Arctica islandica* from the Lillo formation, which was used for seasonality reconstructions (image credit: Doris Smudde). Credit: Doris Smudde

In a warmer climate, summers warm much faster than winters. That is

the conclusion of research into fossil shells by Earth scientist Niels de Winter. With this knowledge, we can better map the consequences of current global warming in the North Sea area.

Alongside colleagues from institutions such as the Institute for Natural Sciences in Brussels, de Winter, affiliated with the Department of Earth Sciences at Vrije Universiteit Amsterdam and the AMGC research group at Vrije Universiteit Brussel, measured the chemical composition of fossil shells from Antwerp, Belgium. Those shells originate from mollusks such as oysters, cockles, and scallops found during the construction works of the Kieldrecht Lock.

The mollusks lived during the Pliocene, approximately three million years ago, in the North Sea, which at that time also covered parts of Flanders and the Netherlands. The shells grew layer by layer, much like [tree rings](#) or fingernails, and stored very detailed information in their [shell](#) during their lifetime.

During the Pliocene, the Earth was on average 2.5 to 3 degrees Celsius warmer than it is now. In their study, [published](#) in *Science Advances*, the researchers took a "snapshot" of the climate at that time to gain insight into the difference between the seasons in a [warmer climate](#).

They use the "clumped isotope analysis" method. With this method, researchers study the composition of shells in even more detail. They do this by measuring the extent to which rare heavy isotopes of both oxygen and carbon occur in the same carbonate from which shells are built. These isotopes are more common in shells that formed in colder waters.

As a result, the measurements can be used to reconstruct the temperature at which the shells were formed. This method is more accurate than conventional methods for temperature reconstructions because it does not rely on assumptions about the composition of the seawater in which

the mollusks grew.

The key insight is that summers warm much more than winters in a warmer climate such as the Pliocene. While winters became about 2.5 degrees warmer, temperatures during summer were about 4.3 degrees higher. The researchers see a similar result in models projecting future climate, which predict roughly the same amount of warming for the year 2100.

The study gives us a glimpse of what the climate in Europe will be like if we continue our current trend toward a warmer world. De Winter said, "We will likely experience stronger temperature differences between summer and [winter](#), and the chance of heat waves during the [summer](#) will increase."

**More information:** Niels de Winter, Amplified seasonality in western Europe in a warmer world, *Science Advances* (2024). [DOI: 10.1126/sciadv.adl6717](#). [www.science.org/doi/10.1126/sciadv.adl6717](http://www.science.org/doi/10.1126/sciadv.adl6717)

Provided by Free University of Brussels

Citation: Summers warm up faster than winters, fossil shells from Antwerp show (2024, May 15) retrieved 11 June 2024 from <https://phys.org/news/2024-05-summers-faster-winters-fossil-shells.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.