

A research team is developing a method to recycle more plastics

May 22 2024



Credit: Unsplash/CC0 Public Domain

Despite consumer efforts to sort and separate recyclables, most plastic bottles still end up in the landfill. Standard recycling methods to sort, shred and remake plastics are limited to just type-1 and type-2

plastics—basically only soda bottles, water bottles and milk jugs.

Global plastic production has increased from 2 million tons in 1950 to 360 million tons in 2018, and about 50% of that plastic becomes trash after a single use. By 2050, it's predicted that 12 billion tons of plastic waste will be in the environment and landfills.

To improve recycling rates, Kevin Schug, the Shimadzu Distinguished Professor of Analytical Chemistry at The University of Texas at Arlington, is working on new ways to separate and recycle mixed plastics. He and a team of graduate and undergraduate researchers at UTA have collaborated on a study [published](#) in *Journal of Chromatography A*.

"A prominent means of chemical recycling is called pyrolysis," Schug said. "During pyrolysis, plastics are heated in an oxygen-free environment until they decompose into pyrolysis oils. These oils have much of the same characteristics as [crude oil](#), with a few exceptions. Importantly, they can be further refined into fuels, and even better, turned into chemical feedstocks to make new plastics."

Unlike traditional plastic recycling that requires sorting and shredding before the material can be recycled, pyrolysis is not limited to specific plastic types. It can accommodate them all.

However, the pyrolysis of mixed plastic waste does create some complex mixtures that manufacturers must examine closely. Contaminants such as sulfur and nitrogen can create chemical compounds that can hurt downstream processing strategies.

"Pyrolysis has become quite a big deal. Many companies are ramping up large chemical [recycling](#) operations," Schug said. "Still, the characterization of the pyrolysis oils requires the development of new

analytical methods, such as the one we describe in our new peer-reviewed research."

With the support of Jean-Francois Borny from Lummus Technologies LLC, a Houston-based chemical company, Schug and his colleagues at UTA—graduate students Alexander Kaplitz and Niray Bhakta, and undergraduate researchers Shane Marshall and Sadid Morshed—created a new supercritical fluid chromatography method that can separate the [pyrolysis](#) oils. The researchers found they could clearly differentiate oils created from polyethylene versus polypropylene feedstocks.

"This is just the beginning, but we're very excited at the potential of this technique to differentiate oils produced from many different plastics and mixtures," Schug said. "Finding a way to better recycle these plastics will help us reduce our reliance on new fossil fuels, and hopefully, do our part to stop contributing to climate change."

More information: Alexander S. Kaplitz et al, Discrimination of plastic waste pyrolysis oil feedstocks using supercritical fluid chromatography, *Journal of Chromatography A* (2024). [DOI: 10.1016/j.chroma.2024.464804](#)

Provided by University of Texas at Arlington

Citation: A research team is developing a method to recycle more plastics (2024, May 22) retrieved 16 June 2024 from <https://phys.org/news/2024-05-team-method-recycle-plastics.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is

provided for information purposes only.