


First use of weather forecasts to show human impact on extreme weather is 'transformational,' scientists say

May 30 2024



An example of a next generation, high resolution and local climate model created by Oxford researchers. Credit: Oxford University Physics

National forecasting centers like the Met Office could apply the same tools used for weather forecasting to quantify how human behavior is

aggravating major events like floods, heat waves and storms, climate scientists at Oxford University Physics show in a [study](#) published today in *Nature Communications*.

Oxford climate physicists, led by Professor Myles Allen, have, for the first time, demonstrated how state-of-the-art weather forecasts can be used to show how greenhouse gas emissions affect [extreme weather](#). In new studies of recent events in both the U.K. and U.S., they assessed the impact of global warming at a local scale and found that human activity both worsened specific weather events and made them more likely to occur.

Their findings coincide with the United Nations "[AI for Good Summit](#)" in Geneva, where scientists from the Oxford Physics team will lead sessions on how [artificial intelligence](#) and machine learning can improve regional forecasting of extreme weather and future climate predictions.


"Weather forecasters could—and should—both warn people of extreme weather and explain how it is being affected by climate change," said Professor Myles Allen, who leads the Oxford University Physics research team. "It isn't a simple case of climate change making all weather worse: some events, like prolonged winter cold, have become less likely."

The new Oxford studies used the world's most reliable medium range weather forecasting model, from the European Center for Medium-Range Weather Forecasting, to assess the impact of climate change on extreme weather.

A [previous study](#), published in *Environmental Research: Climate*, focused on Storm Eunice in the U.K., which reached wind speeds of 122 miles

per hour and caused 17 deaths in February 2022.

"We found that climate change expanded how much of the U.K. was impacted by storm Eunice and intensified the storm's severity by as much as 26%," said Shirin Ermis (Oxford University Physics), who led the U.K. study by Oxford University Physics. The study published today applied the same approach to the U.S. Pacific Northwest heat wave, thought to have killed more than 800 people in June 2021.



A comparison between the limited resolution of current climate models (left) and next generation, higher resolution and more local models (right) created by the Oxford researchers. Credit: Oxford University Physics

"Climate change and human influence is having a very clear impact on certain extreme weather like storms and heat waves," said Dr. Nicholas Leach (Oxford University Physics) who led the U.S. study. "Human influence made this 2021 heat wave at least eight times more likely, and we also found the risk of similar heat waves occurring is doubling every

20 years at the current rate of global warming."

Understanding how climate change and human activity impacts extreme weather events remains a significant and urgent challenge because every year such events cost many lives and billions of dollars in aid and disaster relief around the world.

In the U.K., the cost of dealing with natural disasters caused by extreme weather and climate change could bankrupt the country by the end of the century, according to a recent report from the environmental intelligence agency Kisters. And in the U.S., the cost of dealing with 28 separate weather and climate disasters in 2023 alone topped a record US\$90 billion.

To investigate the impact of climate change on extreme weather, and assess the influence of human activity, scientists rely on computer modeling. However, [climate models](#) are often inaccurate at a regional or local level and only represent specific atmospheric processes at a coarse scale, making their predictions unreliable, especially for extreme weather like storms.

The Oxford teams overcame this by using high-resolution weather forecasting models to simulate extreme weather as if it had occurred in a world without human influence on climate, and in a warmer world of the future. Their models could simulate and predict even unprecedented weather events and can also be used to understand and quantify how human behavior is changing them.

At the AI for Good summit in Geneva today, Professor Philip Stier of Oxford University Physics will convene a workshop with leading international experts to discuss future climate prediction systems. These are expected to make extensive use of artificial intelligence, to deliver more accurate predictions of the impact of [climate change](#) at local level.

More information: Nicholas J. Leach et al, Heatwave attribution based on reliable operational weather forecasts, *Nature Communications* (2024). [DOI: 10.1038/s41467-024-48280-7](https://doi.org/10.1038/s41467-024-48280-7)

Shirin Ermis et al, Event attribution of a midlatitude windstorm using ensemble weather forecasts, *Environmental Research: Climate* (2024). [DOI: 10.1088/2752-5295/ad4200](https://doi.org/10.1088/2752-5295/ad4200)

Provided by University of Oxford

Citation: First use of weather forecasts to show human impact on extreme weather is 'transformational,' scientists say (2024, May 30) retrieved 18 June 2024 from <https://phys.org/news/2024-05-weather-human-impact-extreme-scientists.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.