

New research challenges black holes as dark matter explanation

June 24 2024



Artist's impression of a microlensing event caused by a black hole observed from Earth toward the Large Magellanic Cloud. The light of a background star located in the LMC is bent by a putative primordial black hole (lens) in the Galactic halo and magnified when observed from the Earth. Microlensing causes very characteristic variation of brightness of the background star, enabling the determination of the lens's mass and distance. Credit: J. Skowron / OGLE. Background image of the Large Magellanic Cloud: generated with bsrender written by Kevin Loch, using the ESA/Gaia database



The gravitational wave detectors LIGO and Virgo have detected a population of massive black holes whose origin is one of the biggest mysteries in modern astronomy. According to one hypothesis, these objects may have formed in the very early universe and may include dark matter, a mysterious substance filling the universe.

A team of scientists from the OGLE (Optical Gravitational Lensing Experiment) survey from the Astronomical Observatory of the University of Warsaw have announced the results of nearly 20-year-long observations indicating that such massive black holes may comprise at most a few percent of <u>dark matter</u>. Another explanation, therefore, is needed for gravitational wave sources. The results of the research were <u>published</u> in a study in *Nature* and <u>a study</u> in *The Astrophysical Journal Supplement Series*.

Various astronomical observations indicate that ordinary matter, which we can see or touch, comprises only 5% of the total mass and energy budget of the universe. In the Milky Way, for every 1 kg of ordinary matter in stars, there is 15 kg of dark matter, which does not emit any light and interacts only by means of its gravitational pull.

"The nature of dark matter remains a mystery. Most scientists think it is composed of unknown elementary particles," says Dr. Przemek Mr.óz from the Astronomical Observatory, University of Warsaw, the lead author of both articles. "Unfortunately, despite decades of efforts, no experiment (including experiments carried out with the Large Hadron Collider) has found new particles that could be responsible for dark matter."

Since the first detection of gravitational waves from a merging pair of black holes in 2015, the LIGO and Virgo experiments have detected



more than 90 such events. Astronomers noticed that black holes detected by LIGO and Virgo are typically significantly more massive (20–100 solar masses) than those known previously in the Milky Way (5–20 solar masses).

"Explaining why these two populations of black holes are so different is one of the biggest mysteries of modern astronomy," says Dr. Mr.óz.

One possible explanation postulates that LIGO and Virgo detectors have uncovered a population of primordial black holes that may have formed in the very early universe. Their existence was first proposed over 50 years ago by British theoretical physicist Stephen Hawking, and independently, by the Soviet physicist Yakov Zeldovich.

"We know that the early universe was not ideally homogeneous—small density fluctuations gave rise to current galaxies and galaxy clusters," says Dr. Mr.óz. "Similar density fluctuations, if they exceed a critical density contrast, may collapse and form black holes."

Since the first detection of gravitational waves, more and more scientists have been speculating that such primordial black holes may comprise a significant fraction, if not all, of dark matter.

Fortunately, this hypothesis can be verified with astronomical observations. We observe that copious amounts of dark matter exist in the Milky Way. If it were composed of black holes, we should be able to detect them in our cosmic neighborhood. Is this possible, given that black holes do not emit any detectable light?

According to Einstein's theory of general relativity, light may be bent and deflected in the gravitational field of massive objects, a phenomenon called <u>gravitational microlensing</u>.



"Microlensing occurs when three objects—an observer on Earth, a source of light, and a lens—virtually ideally align in space," says Prof. Andrzej Udalski, the principal investigator of the OGLE survey. "During a <u>microlensing</u> event, the source's light may be deflected and magnified, and we observe a temporary brightening of the source's light."

The duration of the brightening depends on the mass of the lensing object: the higher the mass, the longer the event. Microlensing events by solar mass objects typically last several weeks, whereas those by black holes that are 100 times more massive than the sun would last a few years.

The idea of using gravitational microlensing to study dark matter is not new. It was first proposed in the 1980s by Polish astrophysicist Bohdan Paczyński. His idea inspired the start of three major experiments: Polish OGLE, American MACHO, and French EROS. The first results from these experiments demonstrated that black holes less massive than one solar mass may comprise less than 10% of dark matter. These observations were not, however, sensitive to extremely long-timescale microlensing events, and therefore, not sensitive to massive black holes, similar to those recently detected with gravitational-wave detectors.





Expected vs. observed microlensing events by massive objects toward the Large Magellanic Cloud as seen through the Milky Way halo. If the dark matter in the Universe consisted of putative primordial black holes, over 500 microlensing events would be detected during the OGLE survey in the years 2001-2020. In reality, the OGLE project registered only 13 detections of microlensing events, most likely caused by regular stars. Credit: J. Skowron / OGLE. Background image of the Large Magellanic Cloud: generated with bsrender written by Kevin Loch, using the ESA/Gaia database

In the new article in *The Astrophysical Journal Supplement Series*, OGLE astronomers present the results of nearly 20-year-long photometric monitoring of almost 80 million stars located in a nearby galaxy, called the Large Magellanic Cloud, and the searches for gravitational microlensing events. The analyzed data was collected during the third and fourth phases of the OGLE project from 2001 to 2020.

"This data set provides the longest, largest, and most accurate photometric observations of stars in the Large Magellanic Cloud in the



history of modern astronomy," says Prof. Udalski.

The second article, published in *Nature*, discusses the astrophysical consequences of the findings.

"If the entire dark matter in the Milky Way was composed of black holes of 10 solar masses, we should have detected 258 microlensing events," says Dr. Mr.óz. "For 100 solar mass black holes, we expected 99 microlensing events. For 1,000 solar mass black holes—27 microlensing events."

In contrast, the OGLE astronomers have found only 13 microlensing events. Their detailed analysis demonstrates that all of them can be explained by the known stellar populations in the Milky Way or the Large Magellanic Cloud itself, not by black holes.

"That indicates that <u>massive black holes</u> can compose at most a few percent of dark matter," says Dr. Mr.óz.

The detailed calculations demonstrate that black holes of 10 <u>solar masses</u> may comprise at most 1.2% of dark matter, 100 solar mass black holes—3.0% of dark matter, and 1,000 solar mass black holes—11% of dark matter.

"Our observations indicate that primordial black holes cannot comprise a significant fraction of the dark matter, and simultaneously, explain the observed black hole merger rates measured by LIGO and Virgo," says Prof. Udalski.

Therefore, other explanations are needed for massive <u>black holes</u> detected by LIGO and Virgo. According to one hypothesis, they formed as a product of the evolution of massive, low-metallicity stars. Another possibility involves mergers of less massive objects in dense stellar



environments, such as globular clusters.

"Our results will remain in astronomy textbooks for decades to come," adds Prof. Udalski.

More information: Przemek Mróz et al, No massive black holes in the Milky Way halo, *Nature* (2024). DOI: 10.1038/s41586-024-07704-6. www.nature.com/articles/s41586-024-07704-6. On *arXiv*: DOI: 10.48550/arxiv.2403.02386

Przemek Mróz et al, Microlensing Optical Depth and Event Rate toward the Large Magellanic Cloud Based on 20 yr of OGLE Observations, *The Astrophysical Journal Supplement Series* (2024). DOI: <u>10.3847/1538-4365/ad452e</u>

Provided by University of Warsaw

Citation: New research challenges black holes as dark matter explanation (2024, June 24) retrieved 30 June 2024 from <u>https://phys.org/news/2024-06-black-holes-dark-explanation.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.