


Climate change impacts vary among tree species within the same habitat, finds study

June 14 2024, by Zhang Nannan



Location of the sampling site (red star), meteorological stations (circles), and grid data (crosses). Credit: *Global Ecology and Conservation* (2024). DOI: 10.1016/j.gecco.2024.e02999

Forests, which cover 31% of the Earth's land surface, are essential for maintaining biodiversity and regulating the climate. However, climate change poses a significant challenge to forest growth, with different tree species responding very differently to extreme drought conditions.

As the frequency and intensity of extreme [drought](#) events increase under [global warming](#), there is an urgent need for in-depth research into the climate adaptability of different tree species, in order to take effective measures to protect [forest ecosystems](#) and combat the effects of global warming.

According to a study published in [Global Ecology and Conservation](#), a research team led by Profs. Cai Qiufang and Liu Yu from the Institute of Earth Environment of the Chinese Academy of Sciences (CAS) has investigated the changes in *Pinus tabulaeformis* Carr. and *Tsuga chinensis* Pritz. on the southern slopes of the Qinling Mountains (SSQM) in the context of global warming.

The Qinling Mountains mark the southernmost distribution boundary for natural *P. tabulaeformis* and the northernmost boundary for natural *T. chinensis*.

The researchers found that the climate response patterns of *P. tabulaeformis* and *T. chinensis* on the SSQM have undergone qualitative changes with global warming. The sensitivity of *P. tabulaeformis* to temperature and precipitation has decreased over time, while *T. chinensis* has become more dependent on hydrological conditions.

During extreme drought events, *P. tabulaeformis* shows a stronger drought resistance compared to *T. chinensis*. The stronger drought resistance of *P. tabulaeformis* implies a greater ability to adapt to the warming and drying climate trend, potentially giving it a greater advantage in the current forest ecological composition. In contrast, *T.*

chinensis is less adaptable.

This study highlights that different tree species in the same habitat exhibit significant heterogeneity in growth-climate response, and forest management and [conservation measures](#) should be species-specific.

More information: Mei Xie et al, Assessing climatic response and drought resilience in growth of *Pinus tabulaeformis* Carr. and *Tsuga chinensis* Pritz. on the southern slope of the Qinling Mountains, *Global Ecology and Conservation* (2024). [DOI: 10.1016/j.gecco.2024.e02999](https://doi.org/10.1016/j.gecco.2024.e02999)

Provided by Chinese Academy of Sciences

Citation: Climate change impacts vary among tree species within the same habitat, finds study (2024, June 14) retrieved 18 June 2024 from <https://phys.org/news/2024-06-climate-impacts-vary-tree-species.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.