

## Dark excitons shed new light on matter

June 12 2024



Left: The device comprises a 2D semiconductor between two mirrors, the polariton in purple, and the dark excitons in red or blue depending on spin. The effect of dark excitons can be encapsulated in the polaron, the white dashed

circle. Right: The resulting transmission spectrum. Credit: FLEET

A team of Monash researchers have uncovered for the first time the full effects of interactions between exciton-polaritons and their associated dark excitonic reservoir. The study, "<u>Polaronic polariton quasiparticles</u> in a dark excitonic medium," was published in *Physical Review B*.



Exciton-polaritons (polaritons, for short) are hybrid mixtures of light and matter that inherit the best properties of both. They form in semiconductors sandwiched between two mirrors, through which a laser is shone.

"Polaritons capture useful properties of both matter and light," says lead author Kenneth Choo. "Their 'matter' part means we can manipulate them through their interactions. Meanwhile, the 'light' part makes the <u>polariton</u> almost massless, resulting in the formation of a superfluid condensate that flows effortlessly.

"Put them together, and a new kind of 'liquid light' that we can guide and control is formed, potentially the basis for the next generation of computing that uses light to build circuits," says Kenneth, who is a Ph.D. candidate at Monash.

But with every yang there is a yin, and so it is with the polariton's dark counterpart—a dark exciton.

These dark excitons are uncoupled to light yet still able to interact with the polaritons.

"Whenever the polaritons are formed in experiments, they tend to be accompanied by a substantial <u>reservoir</u> of these dark excitons which form together with them. The fact that they are dark, however, means it has been difficult to see their true effect," says corresponding author Prof Meera Parish.

## A famous technique dresses up for the occasion

The Monash researchers used a framework known as Fermi liquid theory to explain the effect of the dark reservoir on the bright polaritons. The essential idea is to map the (many, many) possible interactions onto



a new 'quasiparticle'—one with different masses, energies and lifetimes to the original. This allows the many-particle system to be treated as if it were a single particle.

One such quasiparticle is the polaron, which can be thought of as a 'zone of influence' around the original particle.

By using this technique and including all possible two-particle interactions, they found that the bright polariton becomes 'dressed' by excitations of the dark medium, forming a new quasiparticle called the polaron polariton.

Physically, this shows up as a reduction in the strength of the lightmatter coupling as the dark exciton density is increased, an effect which is commonly seen in experiment.

In contrast, the strength of the interactions between the dark reservoir and the polaritons are increased, especially when the spins of the reservoir and polariton are different. This effect can be so strong that an additional quasiparticle is generated—the biexciton polariton—which is associated with the bound state of two opposite-spin excitons.

This phenomenon, known as a Feshbach resonance, was first observed in ultracold atomic gases, and offers the promise of being able to tune the interaction strength to whatever value is desired. It would then be possible to create reservoir 'hills' and 'valleys' that guide the flow of the polariton condensate, and then change them around at the flick of a (laser) switch.

## What's next?

This study opens up new opportunities in the field of reservoir engineering. Already, there has been experimental progress in using this



dark reservoir to trap and focus the bright polaritons, and the study shows how the reservoir can be used to fundamentally change the polaritons themselves.

In addition, this study provides a new way to look at old results—by accounting for the hitherto unknown influence of the reservoir, a more accurate determination of interaction strengths can be obtained.

More information: Kenneth Choo et al, Polaronic polariton quasiparticles in a dark excitonic medium, *Physical Review B* (2024). DOI: 10.1103/PhysRevB.109.195432. On *arXiv*: DOI: 10.48550/arxiv.2312.00985

## Provided by FLEET

Citation: Dark excitons shed new light on matter (2024, June 12) retrieved 26 June 2024 from <u>https://phys.org/news/2024-06-dark-excitons.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.