

## Efficient recycling process for rare earth elements through bioleaching and bioaccumulation

June 11 2024



Credit: IMC University of Applied Sciences Krems

A research collaboration between BOKU Tulln and IMC University of Applied Sciences Krems is using the further development of bioleaching



and bioaccumulation to develop a two-stage, environmentally friendly and sustainable process for recovering rare earth elements (rare earths).

In the bioaccumulation step, metal recovery rates of up to 85% were achieved from electronic scrap. The key to success lies in the combination of biotechnological processes. The promising foundations for these methods, which are currently under development, were recently published in *Frontiers in Microbiology*.

The sharp rise in demand for electronics in recent years, used in a wide range of electronic devices such as mobile phones, <u>electric vehicles</u> and computers, has led to an increase in waste containing rare earths. Most of this waste still ends up in landfills unused, despite the fact that rare earths are an important source of raw materials and have even been classified as critical raw materials by the EU.

For this reason, intensive research is being carried out into efficient methods of recovery. Compared to other methods, the microbiologybased methods of bioleaching and bioaccumulation represent a promising green alternative technology for recovering critical raw materials from electronic waste. It is cost-effective, does not produce hazardous or polluting secondary waste, and uses less energy.

The basic principles of the processes are based on the production of acids by certain microorganisms that can leach certain metals such as iron, copper or aluminum from the electronic waste. These metals interfere with the absorption process of valuable rare earths in the subsequent bioaccumulation. Both methods have been researched for some time by the two partners, BOKU Tulln and IMC University of Applied Sciences Krems, and the research teams have now joined forces in a promising collaboration and combined their expertise.

## **'Nothing comes from nothing': Training for microbes**



In addition to the researchers, the current study involved a number of other key players in the bioleaching process, which is summarized in the joint technology: bacteria of different species. For example, Acidithiobacillus thiooxidans and Alicyclobacillus disulfidooxidans, which were originally collected from an acidic mining lake (pH 2.6) in the Czech Republic and then grown together in the laboratory, were used in the bioleaching process. These acidophilic and chemolithotrophic organisms thrive in acidic environments and derive their energy from the oxidation of inorganic compounds.

In terms of bioaccumulation, Escherichia coli, the well-known intestinal bacterium, proved to be the most successful accumulator of rare earths.

The main practical challenge for the enrichment process used to recover rare earths is the high content of other metals typically found in e-waste. In particular, iron, copper and aluminum interfere with the biotechnological process. To overcome this problem, the researchers came up with another innovative option: "training" the microbes. Using a device called a morbidostat developed at IST-Klosterneuburg, the organisms are gradually accustomed to higher metal concentrations. However, the bioaccumulation process has to be carried out carefully so that the organisms do not lose their ability to accumulate the valuable substances.

## **Efficiency in stages**

The methods currently used to extract rare earths are based on <u>chemical</u> <u>processes</u>, which are associated with the formation of environmentally harmful by-products and the creation of new problematic substances. A combination of biotechnological methods has clear advantages over chemical methods, as both the leaching and the accumulation in the cells of the bacteria are environmentally friendly and sustainable, and no



hazardous or polluting substances are produced at any stage of the process.

However, further research is needed to overcome the wide variation in the composition of e-waste. Even if the concentration of interfering metals such as aluminum, iron or copper is changed, the technology must work in such a way that the results are reproducible and reliable.

The researchers at BOKU and IMC Krems are pursuing several strategies to achieve this. Another strategy is to acclimate the bacteria responsible for bioleaching and bioaccumulation to high concentrations of interfering metals. This is made possible by using a system called morbidostat. In this system, the micro-organisms are exposed to a gradually increasing concentration of interfering metals. Then researchers wait until acculturation occurs and the organisms start to grow further.

Together with the conditioning of the microorganisms, systems are being tested that can trigger a reduction in the concentration of interfering metals. The materials being investigated include the so-called lignin hydrogels developed at BOKU. The combination of these strategies aims to ensure the efficiency and sustainability of the innovative combination of bioleaching and <u>bioaccumulation</u> in order to develop a new, environmentally-friendly method for recycling scarce <u>rare earths</u>.

## Provided by IMC University of Applied Sciences Krems

Citation: Efficient recycling process for rare earth elements through bioleaching and bioaccumulation (2024, June 11) retrieved 26 June 2024 from <a href="https://phys.org/news/2024-06-efficient-recycling-rare-earth-elements.html">https://phys.org/news/2024-06-efficient-recycling-rare-earth-elements.html</a>

This document is subject to copyright. Apart from any fair dealing for the purpose of private



study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.