

## A fierce battle is being fought in the soil beneath our feet—and the implications for global warming could be huge

June 6 2024, by Kristine Crous, Belinda Medlyn and David S Ellsworth



The ecosystem P budget. The ecosystem P budget under ambient CO2 (aCO2) (a) and eCO2 (b) treatment, assembled from data collected at EucFACE over 6 years (2013–2018). Credit: *Nature* (2024). DOI: 10.1038/s41586-024-07491-0

As humanity continues to burn fossil fuels, the delicate balance of life on Earth is changing. That's true of trees, many of which are growing faster as a result of increased carbon dioxide  $(CO_2)$  concentrations in our atmosphere.



But not all trees are responding in this way. In particular, eucalypts—Australia's iconic forest trees—haven't benefited from the increase in  $CO_2$  as they were expected to.

Why not? <u>Our new research</u>, published today in *Nature*, shows it comes down to a below-ground battle for <u>phosphorus</u>, a mineral nutrient in soils that is essential for <u>tree growth</u>. The results suggest in some parts of the world, increased  $CO_2$  means tiny bugs in the soil "hold onto" their phosphorus, making less available for trees.

This is alarming news for some forests, because according to current projections, global forest growth is meant to limit damage from <u>global</u> <u>warming</u>.

## What our study involved

Our study used data from a Western Sydney University experiment known as "Eucalyptus Free Air  $CO_2$  Enrichment," or <u>EucFACE</u>. The experiment is located in a century-old Cumberland plain woodland in Sydney's Hawkesbury district.

 $CO_2$  is released into the woodland through a computer-controlled system. Scientists then monitor the effects on trees, soils and the broader ecosystem. Over six years,  $CO_2$  was raised to the levels expected around the year 2050 (according to the current business-as-usual emissions trajectory).

<u>Our previous studies</u> found the woodland trees did not show any extra growth at high  $CO_2$  levels. We suspected the low availability of soil phosphorus was the cause, and set out to test this.

Phosphorus is crucial to the process of photosynthesis that makes trees grow. Phosphorus in soil is provided by bugs known as microbes. These



micro-organisms break down dead and decaying matter, and in the process change phosphorus into a form that plants can take up with their roots.

Most Australian soils are naturally low in phosphorus, because they are derived from ancient, nutrient-depleted rocks. The same is true for most soils in tropical and subtropical regions. That makes the phosphorus service provided by microbes even more important.

We sampled phosphorus in all parts of the ecosystem, tracing its journey from the soil to the trees. We found under high- $CO_2$  conditions the microbes keep more of the phosphorus they produce, to aid their own metabolism. This left less available for trees to take up.

This occurred despite the trees trying to "bargain" for phosphorus by releasing extra carbon into the soil to feed the microbes.

What's more, trees are big "recyclers" of phosphorus—they remove half of the phosphorus from any leaf before it falls. But this was still not enough to support extra tree growth.

## Why this matters

Our study is the first to show how the phosphorus cycle is affected by high  $CO_2$ —and in particular, the role of soil microbes.

The results are important to predicting <u>soil</u> phosphorus availability, and plant productivity, in woodlands and forests as  $CO_2$  levels increase in the atmosphere.

Current climate projections assume increasing  $CO_2$  will lead to <u>more</u> <u>forest growth globally</u>. Forests are a vital carbon "sink"—that is, they draw down carbon from the atmosphere. So the increased forest growth



was projected to go some way to limiting the effects of climate change.

If our results are taken into account, future warming would be higher than current projections. However, it's important to verify our results in other locations, with other tree species. New experiments are being formed by overseas teams, including in the Amazon rainforest, to test the findings.

Importantly, our results don't mean that forests are not a crucial sink for carbon. Forests hold a vast quantity of carbon. Avoiding deforestation and planting new forests are both valuable means of maintaining and adding to carbon stores.

Our research demonstrates the importance of considering soils when growing trees. We also hope our research stimulates further efforts to find phosphorus in ecosystems, especially in <u>tropical rainforests</u> where phosphorus is often greatly limited.

**More information:** Mingkai Jiang et al, Microbial competition for phosphorus limits the  $CO_2$  response of a mature forest, *Nature* (2024). DOI: 10.1038/s41586-024-07491-0

This article is republished from <u>The Conversation</u> under a Creative Commons license. Read the <u>original article</u>.

Provided by The Conversation

Citation: A fierce battle is being fought in the soil beneath our feet—and the implications for



global warming could be huge (2024, June 6) retrieved 26 June 2024 from <u>https://phys.org/news/2024-06-fierce-fought-soil-beneath-feet.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.