

# CO<sub>2</sub> puts heavier stamp on temperature than previously thought, analysis suggests

June 24 2024

---



Credit: Pixabay/CC0 Public Domain

A doubling of the amount of CO<sub>2</sub> in the atmosphere could cause an increase in the average temperature on Earth from 7 to a maximum of 14 degrees. This is shown in the analysis of sediments from the Pacific Ocean off the coast of California, by researchers at NIOZ and the Universities of Utrecht and Bristol. Their results were [published](#) in *Nature Communications*.

"The temperature rise we found is much larger than the 2.3 to 4.5

degrees that the UN climate panel, IPCC, has been estimating so far," said the first author, Caitlyn Witkowski.

The researchers used a 45-year-old drill core extracted from the bottom of the Pacific Ocean. "I realized that this core is very attractive for researchers, because the [ocean floor](#) at that spot has had oxygen-free conditions for many millions of years," said Professor Jaap Sinninghe Damsté, senior scientist at NIOZ and professor of organic geochemistry at Utrecht University.

"As a result, [organic matter](#) is not broken down as quickly by microbes and more carbon is preserved," Damsté said. He was also the supervisor of Witkowski, whose doctorate thesis included this research.

"CO<sub>2</sub> over the past 15 million years has never before been examined from a single location," Witkowski said. The upper thousand meters of the drill core correspond to the past 18 million years. From this record, the researchers were able to extract an indication of the past seawater temperature and an indication of ancient atmospheric CO<sub>2</sub> levels, using a new approach.

## **Derived temperature**

The researchers derived the temperature using a method developed 20 years ago at NIOZ, called the TEX<sub>86</sub> method. "That method uses specific substances that are present in the membrane of archaea, a distinct class of microorganisms," Damsté explains.

"Those archaea optimize the chemical composition of their membrane depending on the temperature of the water in the upper 200 meters of the ocean. Substances from that membrane can be found as molecular fossils in the ocean sediments, and analyzed to this day."

## CO<sub>2</sub> from chlorophyll and cholesterol

The researchers developed a new approach to derive past atmospheric CO<sub>2</sub> content by using the [chemical composition](#) of two specific substances commonly found in algae: chlorophyll and cholesterol. This is the first study to use cholesterol for quantitative CO<sub>2</sub> and the first study to use chlorophyll for this time period. To create these substances, algae must absorb CO<sub>2</sub> from the water and fix it via photosynthesis.

Damsté said, "A very small fraction of the carbon on Earth occurs in a 'heavy form,' <sup>13</sup>C instead of the usual <sup>12</sup>C. Algae have a clear preference for <sup>12</sup>C. However, the lower the CO<sub>2</sub> concentration in the water, the more algae will also use the rare <sup>13</sup>C. Thus, the <sup>13</sup>C content of these two substances is a measure of the CO<sub>2</sub> content of the ocean water. And that in turn, according to solubility laws, correlates with the CO<sub>2</sub> content of the atmosphere."

Using this new method, it appears that the CO<sub>2</sub> concentration dropped from about 650 parts per million, 15 million years back, to 280 just before the industrial revolution.

## Stronger relationship

When the researchers plot the derived temperature and atmospheric CO<sub>2</sub> levels of the past 15 million years against each other, they find a strong relationship.

The average temperature 15 million years back was over 18 degrees: 4 degrees warmer than today and about the level that the UN climate panel, IPCC, predicts for the year 2100 in the most extreme scenario.

"So, this research gives us a glimpse of what the future could hold if we take too few measures to reduce CO<sub>2</sub> emissions and also implement few

technological innovations to offset emissions," Damsté said.

"The clear warning from this research is CO<sub>2</sub> concentration is likely to have a stronger impact on [temperature](#) than we are currently taking into account."

**More information:** Caitlyn R. Witkowski et al, Continuous sterane and phytane  $\delta^{13}\text{C}$  record reveals a substantial pCO<sub>2</sub> decline since the mid-Miocene, *Nature Communications* (2024). [DOI: 10.1038/s41467-024-47676-9](#)

Provided by Royal Netherlands Institute for Sea Research

Citation: CO<sub>2</sub> puts heavier stamp on temperature than previously thought, analysis suggests (2024, June 24) retrieved 29 June 2024 from <https://phys.org/news/2024-06-heavier-temperature-previously-thought-analysis.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.