

Iceland's volcano eruptions may last decades, researchers find

June 26 2024



Volcanologists and geochemists getting ready to sample lava during the July 2023 Fagradalsfjall eruption on the Reykjanes Peninsula of Iceland. Credit: Valentin Troll

Iceland's ongoing volcanic eruptions may continue on and off for years



to decades, threatening the country's most densely populated region and vital infrastructure, researchers predict from local earthquake and geochemical data.

The eruptions on the Reykjanes Peninsula have forced authorities to declare a state of emergency, with a series of eight eruptions having occurred since 2021. This southwestern region is home to 70% of the country's population, its only international airport, and several geothermal power plants that supply hot water and electricity. The most recent eruption from May through June triggered the evacuation of residents and visitors from the Blue Lagoon geothermal spa, a popular tourist attraction, for the third time in more than two months.

Although Iceland sees regular eruptions because it sits above a volcanic hot spot, the Reykjanes Peninsula has been dormant for 800 years. Its last volcanic era continued <u>over centuries</u>, however, prompting scientists to predict the renewed volcanism to be the start of a long episode.

Under an hour's drive from the island's capital city Reykjavík, the eruptions pose considerable risks for economic disruption, and they leave evacuated communities uncertain of a possible return.

An international team of scientists has been watching the volcanoes over the past three years. Analyzing seismic tomography imaging and the composition of lava samples, they've uncovered parts of the geological processes behind the new volcanic era. They predict the region may have to prepare for recurring eruptions lasting years to decades and possibly centuries.





Lava from the summer 2023 Fagradalsfjall eruption flowing on top of lava from the summer 2021 eruption. Volcanism has been ongoing since 2021 in the Reykjanes Peninsula, which hosts the majority of Iceland's population. Credit: Valentin Troll

The researchers <u>report their findings</u> in a paper published June 26 in the journal *Terra Nova*. The project included collaborations from the University of Oregon, Uppsala University in Sweden, University of Iceland, Czech Academy of Sciences and University of California, San Diego. The work follows an <u>earlier study</u> in *Nature Communications* of the initial Reykjanes eruptions in 2021.




Almost all of Iceland's island is built from lava, said Ilya Bindeman, a volcanologist and earth sciences professor at the UO. The country sits on the Mid-Atlantic Ridge, the <u>tectonic plate boundary</u> that causes North America and Eurasia to push further apart. The drifting of these plates can spark <u>volcanic eruptions</u> when hot rock from the earth's mantle—the middle and largest layer of the planet—melts and rises to the surface.

Although scientists know the origin of Reykjanes Peninsula's current eruptions is plate movement, the kind of <u>magma</u> storage and plumbing systems that feed them are unidentified, Bindeman said. The peninsula consists of eight volcanically active sites, so understanding whether there is one shared magma source or multiple independent ones and their depth can help predict the duration and impact of these eruptions.

Using geochemical and seismic data, the researchers investigated whether the magma of the initial eruptions from one volcano in the peninsula from 2021 to 2023 came from the same source as the magma in the recent eruptions of a different volcano to the west.

Bindeman specializes in isotopic analysis, which can help identify the "fingerprint" of magma. Magma is made of mostly eight elements, including oxygen and hydrogen, and 50 different trace elements in smaller concentrations and various ratios. The unique combination of trace elements can help differentiate magma sources from one another. Scientists can also measure the abundance of isotopes, elements with the same chemical properties but different masses, in the magma. There are three different isotopes of oxygen, for example, Bindeman said.





The equipment used to identify the "fingerprint" of magma sources. Called a laser fluorination line, the apparatus extracts and measures oxygen isotopes. This is housed in volcanologist Ilya Bindeman's lab at the University of Oregon. Credit: Charlie Litchfield/University of Oregon.

"In the air we breathe, there's a mixture of these oxygen isotopes and we don't feel the difference," he said. "Their differences are usually not important for chemical reactions but are important to recognize as their relative abundances in magma can differentiate one magma source from another."

Analyzing samples of lava rock from two different volcanoes on the



peninsula, their similar fingerprints implied a shared magma storage zone below the peninsula. Imaging of Earth's interior based on local earthquakes also suggested the existence of a reservoir about 5.5 to 7.5 miles in Earth's crust, the shallowest layer.

However, that storage is ultimately fed by the melting rock deeper in the mantle, which can cause eruptions that last decades, with hundreds of square miles of magma surfacing, Bindeman said. Iceland's hotspot also will have no problem fountaining that flow, he said.

Although this marks the beginning of potentially persistent volcanic episodes in Iceland, the researchers can't precisely predict yet how long the episodes and the gaps between each will last.

"Nature is never regular," Bindeman said. "We don't know how long and how frequently it will continue for the next ten or even hundred years. A pattern will emerge, but nature always has exceptions and irregularities."





Lava from the January 2024 Sundhnúkur eruption at the outskirts of Grindavik town. Photographed in April 2024. Volcanism has been ongoing since 2021 in the Reykjanes Peninsula, which hosts the majority of Iceland's population. Credit: Valentin Troll

Discussions are continuing on plans to safely drill into the volcanic sites to glean insights into the geological processes driving the eruptions.

Because the <u>volcanic activity</u> is less volatile and explosive than eruptions in other countries, it provides a rare opportunity for scientists to approach fissures actively erupting lava, Bindeman said. He called it a "natural laboratory," both astonishing and chilling.



"When you witness a volcanic eruption, you can feel that these are the massive forces of nature, and you yourself are very small," Bindeman said. "These events are ordinary on the geological scale, but on the human scale, they can be devastating."

More information: The Fagradalsfjall and Sundhnúkur Fires of 2021-2024: a single magma reservoir under the Reykjanes Peninsula, Iceland?, *Terra Nova* (2024). DOI: 10.1111/ter.12733

Provided by University of Oregon

Citation: Iceland's volcano eruptions may last decades, researchers find (2024, June 26) retrieved 17 July 2024 from <u>https://phys.org/news/2024-06-iceland-volcano-eruptions-decades.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.