

Observing flows at a liquid-liquid-solid intersection

June 26 2024, by Rachel Berkowitz





Cameras and lasers measure the flow fields near a liquid-liquid moving contact line. Credit: *The European Physical Journal Special Topics* (2024). DOI: 10.1140/epjs/s11734-024-01170-x

Most of us are familiar with the classic example of a liquid-gas moving contact line on a solid surface: a raindrop, sheared by the wind, creeps along a glass windscreen. The contact line's movements depend on the interplay between viscous and surface tension forces—a relationship that has been thoroughly investigated in experimental fluid mechanics.

In a study <u>published</u> in *The European Physical Journal Special Topics*, Harish Dixit, of the Indian Institute of Technology Hyderabad, and his colleagues now examine the movements of a contact line formed at the interface between two immiscible liquids and a solid. The experiments fill a gap in <u>fluid dynamics</u> and suggest a mechanism for an imposed boundary condition that eludes mathematical description.

According to theory, the movement of a liquid-liquid contact line should be governed entirely by the liquids' viscosity ratio and the angle at which the liquid interface meets the solid. To examine this in a real-world system, Dixit and his colleagues filled a rectangular tank with two liquid layers—silicone oil atop sugar water—with similar densities but significantly different viscosities. The researchers placed a glass slide at the edge of the tank, which they could slide vertically to create a moving contact line.

Using a technique that tracks <u>tiny particles</u> introduced to the liquids and illuminated with <u>laser light</u>, the researchers simultaneously mapped the flow field on both sides of the liquid-liquid interface while they moved the glass slide. They found that the flow velocities rapidly decreased



close to the contact line. Additionally, the liquid interface appeared to slip along the moving glass slide, rather than staying pinned to it—resolving the apparent "singularity" in models that impose a no-slip boundary condition on the moving wall.

The findings lend support to theoretical models of contact line dynamics, and offer parameters that could improve <u>numerical simulations</u> which are based on similar-viscosity liquids.

More information: Charul Gupta et al, An experimental investigation of flow fields near a liquid–liquid moving contact line, *The European Physical Journal Special Topics* (2024). DOI: 10.1140/epjs/s11734-024-01170-x

Provided by SciencePOD

Citation: Observing flows at a liquid-liquid-solid intersection (2024, June 26) retrieved 17 July 2024 from <u>https://phys.org/news/2024-06-liquid-solid-intersection.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.