

Marine CO₂ removal technologies could depend on the appetite of the ocean's tiniest animals

June 17 2024, by Tyler Rohr, Ali Mashayek and Sophie Meyjes



Credit: ESA, CC BY-SA

As the world struggles to decarbonize, it's becoming increasingly clear we'll need to both rapidly reduce emissions and actively remove carbon dioxide (CO_2) from the atmosphere. The latest Intergovernmental Panel on Climate Change <u>report</u> considered 230 pathways to keep global warming below 1.5°C. All require CO_2 removal.



Some of the most promising CO_2 removal technologies receiving government funding in the <u>United States</u>, <u>United Kingdom</u> and <u>Australia</u> seek to increase the massive carbon storage potential of the ocean. These include <u>fertilizing tiny plants</u> and <u>tweaking ocean chemistry</u>.

Ocean-based <u>approaches</u> are gaining popularity because they could potentially store carbon for <u>a tenth of the cost</u> of "<u>direct air capture</u>", where CO_2 is sucked from the air using energy-intensive machinery.

But the marine carbon cycle is much harder to predict. Scientists must unravel the many complex natural processes that could alter the efficiency, efficacy and safety of ocean-based CO_2 removal before it can go ahead.

In <u>our new research</u> published in *Global Biogeochemical Cycles*, we highlight a surprisingly important mechanism that had previously been overlooked. If CO_2 removal techniques change the appetite of tiny animals at the base of the food chain, that could dramatically change how much carbon is actually stored.

Plankton dominate the carbon pump

Tiny marine life forms called <u>plankton</u> play a huge role in ocean carbon cycling. These microscopic organisms drift on the <u>ocean currents</u>, moving captured carbon throughout the seas.

Like plants on land, <u>phytoplankton</u> use sunlight and CO_2 to grow through photosynthesis.

Zooplankton, on the other hand, are tiny animals that mostly eat phytoplankton. They come in many <u>shapes and sizes</u>. If you put them in a lineup, you might think they came from different planets.





A lineup of zooplankton: tiny marine animals that look like they come from different planets. Credit: Julian Uribe-Palomino/IMOS-CSIRO

Across all this diversity, <u>zooplankton</u> have very different <u>appetites</u>. The hungrier they are, they faster they eat.

Uneaten phytoplankton—and zooplankton poo—can sink to great depths, keeping carbon locked away from the atmosphere for centuries. Some even sink to the seafloor, eventually transforming <u>into fossil fuels</u>.

This transfer of carbon from the atmosphere to the ocean is known as the "biological pump." It keeps hundreds of billions of tons of carbon in the ocean and out of the atmosphere. That translates to about 400 ppm CO₂ and 5° C of cooling.



Picky eaters

In our <u>new research</u>, we wanted to better understand how zooplankton appetites influence the biological pump.

<u>First</u> we had to work out how zooplankton appetites differ across the ocean.

We used a <u>computer model</u> to simulate the seasonal cycle of phytoplankton population growth. This is based on the balance of reproduction and death. The model simulates reproduction really well.

Zooplankton appetites <u>largely determine the death rate</u>. But the model's not so good at <u>simulating death rates</u>, because it doesn't have enough information about zooplankton appetites.

So we tested dozens of different appetites and then checked our results against real-world data.





In the ocean carbon cycle, the biological pump begins with the capture of atmospheric carbon dioxide during photosynthesis by phytoplankton. If the phytoplankton die, the carbon in their bodies is stored deep in the ocean. However, zooplankton grazing will release carbon dioxide back into the atmosphere. Credit: IAEA

To get global observations of phytoplankton seasonal cycles without a fleet of ships, we used satellite data. This is possible even though phytoplankton are tiny, because their light-catching pigments are <u>visible</u> from space.

We ran the model in more than 30,000 locations and found zooplankton appetites vary enormously. That means all those different types of zooplankton are not spread evenly across the ocean. They appear to gather around their favorite types of prey.

<u>In our latest research</u>, we show how this diversity influences the biological pump.

We compared two models, one with just two types of zooplankton and another with an unlimited number of zooplankton—each with different appetites, all individually tuned to their unique environment.

We found including realistic zooplankton diversity reduced the strength of the biological pump by a billion tons of carbon every year. That's bad for humanity, because most of the carbon that doesn't go into the ocean ends up back in the atmosphere.



Not all of the carbon in the bodies of the phytoplankton would have sunk deep enough to <u>be locked away</u> from the atmosphere. But even if only a quarter did, once <u>converted to CO_2 </u>, that could match annual emissions from the entire <u>aviation industry</u>.

The ocean as a sponge

Many ocean-based CO_2 removal technologies will alter the composition and abundance of phytoplankton.

Biological ocean-based CO_2 removal technologies such as "<u>ocean iron</u> <u>fertilization</u>" seek to increase phytoplankton growth. It's a bit like spreading fertilizer in your garden, but on a much bigger scale—with a fleet of ships seeding iron across the ocean.

The goal is to remove CO_2 from the atmosphere and pump it into the deep ocean. However, because <u>some phytoplankton crave iron more than</u> <u>others</u>, feeding them iron could change the composition of the population.

Alternatively, non-biological ocean-based CO_2 removal technologies such as "<u>ocean alkalinity enhancement</u>" shift the chemical balance, allowing more CO_2 to dissolve in the water before it reaches chemical equilibrium. However, the most accessible sources of alkalinity are minerals, including nutrients that encourage the growth of certain <u>phytoplankton over others</u>.

If these changes to phytoplankton favor different types of zooplankton with different sized appetites, they are likely to change the strength of the biological pump. This could compromise—or complement—the efficiency of ocean-based CO_2 removal technologies.



Moving forward from a sea of uncertainty

Emerging private-sector CO_2 removal companies will require accreditation from reliable <u>carbon offset registries</u>. This means they must demonstrate their technology can:

- 1. Remove carbon for hundreds of years (permanence)
- 2. Avoid major environmental impacts (safety)
- 3. Be amenable to accurate monitoring (verification).

Cast against a <u>sea of uncertainty</u>, the time is now for oceanographers to <u>establish the necessary standards</u>.

Our research shows CO_2 removal technologies that change phytoplankton communities could also drive changes in carbon storage, by modifying zooplankton appetites. We need to better understand this before we can accurately predict how well these technologies will work and how we must monitor them.

This will require tremendous effort to overcome the challenges of <u>observing</u>, <u>modeling</u> and <u>predicting</u> zooplankton dynamics. But the payoff is huge. A more reliable regulatory framework could pave the way for a <u>trillion-dollar</u>, morally imperative, emerging CO_2 removal industry.

More information: S. A. Meyjes et al, Impact of Spatial Variability in Zooplankton Grazing Rates on Carbon Export Flux, *Global Biogeochemical Cycles* (2024). DOI: 10.1029/2023GB008085

This article is republished from <u>The Conversation</u> under a Creative Commons license. Read the <u>original article</u>.



Provided by The Conversation

Citation: Marine CO₂ removal technologies could depend on the appetite of the ocean's tiniest animals (2024, June 17) retrieved 29 June 2024 from <u>https://phys.org/news/2024-06-marine-technologies-appetite-ocean-tiniest.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.