

First map of outflows from nearby quasar I Zwicky 1

June 11 2024



Artist's impression of outflows from a quasar. Credit: ESO/M. Kornmesser

SRON astronomers have for the first time mapped the outflows from one of the closest quasars to Earth. Quasars are bright cores of galaxies powered by the supermassive black hole in their center. The team has

probed gas outflows in I Zwicky 1, a close-by quasar, to map its system of clouds being blown away at tens to thousands of kilometers per second. Their findings are published in the journal *Astronomy & Astrophysics*.

Most galaxies, including our Milky Way, harbor a [supermassive black hole](#) in their center. These typically weigh millions of [solar masses](#). Many of them keep lurking in the blackness of space, with little to give them away. Some have, however, large deposits of material in their vicinity to feed on. This turns the central region into a brilliant beacon, outshining the entire host galaxy.

Given their compact size and large distance from Earth, these [active galactic nuclei](#) appear as bright dots, like the Milky Way stars. This is also why they were historically classified as quasi-stellar objects (quasars).

Most quasars reside in the distant, [early universe](#), but I Zwicky 1 is relatively close at less than a billion lightyears away from Earth. This provides astronomers with a convenient laboratory for studying the [extreme conditions](#) in quasars.

A team of astronomers led by Anna Juráňová (SRON), including Elisa Costantini (SRON), has now for the first time mapped its outflows. Using the Hubble Space Telescope, they recovered the properties of four outflows of ionized gas clouds, being blown out at speeds of 60, 280, 1950 and 2900 kilometers per second.

"I Zwicky 1 is very special in its properties," says Juráňová. "Other quasars have similar outflows, but in this one, everything is just right. Our [viewing angle](#), the width of the lines in the spectrum, and so on. This allows us to dig much deeper into its processes. We have created a global picture of the motions of the ionized gas in a quasar, which is

rare."

The team found that one of the outflows was trapped in the shadow of the other one. This results from the strong radiation from the quasar pushing the clouds out and away from the vicinity of the black hole. Ions of elements such as nitrogen, oxygen and carbon within the gas clouds absorb the quasar's ultraviolet light and get pushed away as a consequence. I Zwicky 1 is the nearest quasar offering concrete evidence of this mechanism at play.

The environment around I Zwicky 1 appears more dynamical than what astronomers often see around nearby supermassive black holes. Juráňová says, "Our data suggest that far more gas is being lifted and blown out from the disk around the black hole. Having this insight brings us closer to unraveling the way these supermassive black holes grow and interact with their surroundings."

More information: *Astronomy & Astrophysics* (2024).

Provided by SRON Netherlands Institute for Space Research

Citation: First map of outflows from nearby quasar I Zwicky 1 (2024, June 11) retrieved 18 June 2024 from <https://phys.org/news/2024-06-outflows-nearby-quasar-zwicky.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.