

# Determining refractive index at relativistic speeds

June 26 2024, by Clare Sansom

---



A ray of light refracted through a glass slab. Credit: ajistai

If you studied advanced physics at high school, there's a good chance that you remember Snell's law, which states how a ray of light bends when it crosses a boundary between two media. According to this law,

the ratio of the sines of the incident and refracted angles is a non-universal constant, later understood as the relative refractive index of the refracting medium with respect to the incident one.

The theoretical formulation of the law of optical refraction was initiated by Descartes in the early 17th century. He introduced a metaphysical form of an optical-mechanical analogy, suggesting a conceptual parallel between trajectories in [classical mechanics](#) and wavefronts in optics.

Descartes considered the conservation of the tangential component of the velocity of a particle, as well as light, as it crosses a boundary between [media](#). He determined the relative refractive index as the ratio of the speed of light in the two media. Later this century, Descartes' metaphysical theory was rejected by Fermat by introducing the "least time principle" that was named after him. He also determined the relative refractive index as the ratio of the speed of light in the two media.

Descartes' idea, which pre-dates the discovery that the speed of light is finite was "brilliant." It needed a relativistic theory that was formulated much later in the early 20th century. Shyamal Biswas and his research group members at the University of Hyderabad in India have now extended Descartes' metaphysical theory to determine the mechanical refractive index for relativistic particles entering media at all possible speeds. This work has now been [published](#) in the journal *The European Physical Journal D*.

Biswas and his co-workers defined the mechanical refractive index by comparing the Helmholtz wave equation for light in an optical medium and the time-independent Klein-Gordon equation for a relativistic particle in a potential. "Our [analytical method](#) involved only quite simple mathematics: algebra, trigonometry, and calculus," he explains.

The researchers' calculations of the mechanical refractive [index](#) for particles traveling at different speeds up to that of [light](#) exactly matches with Descartes' result at the non-relativistic limit and Fermat's result at the ultra-relativistic limit. "Our calculations may have [applications](#) in studying the transmission and reflection of particles in mechanical media," concludes Biswas.

**More information:** Bikram Keshari Behera et al, Refractive index for the mechanical refraction of a relativistic particle, *The European Physical Journal D* (2024). [DOI: 10.1140/epjd/s10053-024-00849-z](https://doi.org/10.1140/epjd/s10053-024-00849-z)

Provided by SciencePOD

Citation: Determining refractive index at relativistic speeds (2024, June 26) retrieved 29 June 2024 from <https://phys.org/news/2024-06-refractive-index-relativistic.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.