


# Tackling the dual threat: A global strategy for PM<sub>2.5</sub> and O<sub>3</sub> pollution

June 28 2024



Graphical abstract. Credit: *Eco-Environment & Health* (2024). DOI: 10.1016/j.eehl.2024.04.004

Air pollution is a severe risk to human health and the environment, particularly from fine particulate matter (PM<sub>2.5</sub>) and ozone (O<sub>3</sub>). Despite global efforts, many cities continue to face significant exposure risks from these pollutants.

PM<sub>2.5</sub> and O<sub>3</sub> originate from similar sources and interact in complex ways, compounding their harmful effects. Addressing these intertwined

pollutants requires innovative strategies. Due to these challenges, it is necessary to conduct in-depth research to develop effective strategies for joint PM<sub>2.5</sub> and O<sub>3</sub> control.

A research team from Hubei University of Economics, Nanjing University, and Yangtze University conducted a study on the spatial and temporal patterns of PM<sub>2.5</sub>-O<sub>3</sub> compound pollution. Published in *Eco-Environment & Health*, on 19 April 2024, the research analyzed data from 120 cities worldwide between 2019 and 2022, proposing a framework for synergistic pollution control. The research is [published](#) in the journal *Eco-Environment & Health*.

The study revealed that nearly 50% of cities worldwide are affected by PM<sub>2.5</sub>-O<sub>3</sub> compound pollution, with hotspots in China, Korea, Japan, and India. Significant spatial correlations between PM<sub>2.5</sub> and O<sub>3</sub> concentrations were identified, driven by common precursors such as [nitrogen oxides](#) (NO<sub>x</sub>) and Volatile Organic Compounds (VOCs).

The analysis showed that 52.5% of the cities achieved coordinated reductions in PM<sub>2.5</sub> and O<sub>3</sub>, with average decreases of 13.97% and 19.18%, respectively. This highlights the potential for joint pollution control measures. The proposed framework aims to manage emissions from both pollutants simultaneously, leveraging their spatial and chemical interactions.

Key findings included the identification of cities with high exposure risks and the demonstration of a positive spatial correlation between PM<sub>2.5</sub> and O<sub>3</sub> concentrations, suggesting that integrated control strategies could significantly enhance urban air quality and public health.

Dr. Chao He, lead author of the study, stated, "Our findings underscore the critical need for integrated pollution control strategies. By addressing PM<sub>2.5</sub> and O<sub>3</sub> together, we can more effectively reduce the health risks

and environmental impacts associated with these pollutants."

The proposed synergistic control framework offers a promising approach to managing global [air pollution](#). Implementing these strategies could lead to significant improvements in urban air quality, reducing [health risks](#) and promoting sustainable development. Policymakers and environmental agencies can use these insights to develop more effective regulations and interventions for cleaner, healthier cities.

**More information:** Chao He et al, Synergistic PM<sub>2.5</sub> and O<sub>3</sub> control to address the emerging global PM<sub>2.5</sub>-O<sub>3</sub> compound pollution challenges, *Eco-Environment & Health* (2024). [DOI: 10.1016/j.eehl.2024.04.004](#)

Provided by Chinese Academy of Sciences

Citation: Tackling the dual threat: A global strategy for PM<sub>2.5</sub> and O<sub>3</sub> pollution (2024, June 28) retrieved 8 September 2024 from <https://phys.org/news/2024-06-tackling-dual-threat-global-strategy.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.