


# Untangling the entangled: Quantum study shines fresh light on how neutrinos fuel supernovae

June 22 2024, by Matt Lakin



Flavor inversion probabilities for  $N=12$  neutrinos. The lines show the single-step Trotter (continuous) and exact (dashed) simulations, and the points show the results from H1-1 (dark circles, using 240 shots) and H1-1E (light squares, using 1200 shots). Credit: *Physical Review Letters* (2023). DOI: 10.1103/PhysRevLett.130.221003

Researchers used quantum simulations to obtain new insights into the nature of neutrinos—the mysterious subatomic particles that abound throughout the universe—and their role in the deaths of massive stars.

The [study](#) relied on support from the Quantum Computing User Program, or QCUP, and the Quantum Science Center, a national Quantum Information Science Research Center, at the Department of Energy's Oak Ridge National Laboratory. The work is published in the journal *Physical Review Letters*.

"This understanding was something new that hasn't come out of classical computing systems," said Martin Savage, the study's senior author and a professor of physics at the University of Washington.

"We recognized for the first time we could study how entanglement between multiple neutrinos is induced over time, and these results are within the error bars of what we'd expect from a classical computer. It's a step in the direction of better, more accurate and more scalable quantum simulations."

Neutrinos result from [nuclear reactions](#)—from the huge reactions that cause the sun to shine, to the tiny reactions that enable radioactive tracers for medical tests. These extremely light particles appear everywhere, carry no electric charge and seldom interact with other matter.

But during the collapse and explosion of a star—a process better known as a supernova—neutrinos exchange energy and momentum with not just each other but with everything around them.

"At this point, the neutrinos go from passive particles—almost bystanders—to major elements that help drive the collapse," Savage said. "Supernovae are interesting for a variety of reasons, including as

sites that produce heavy elements such as gold and iron. If we can better understand neutrinos and their role in the star's collapse, then we can better determine and predict the rate of events such as a supernova."

Scientists seldom observe a supernova close-up, but researchers have used classical supercomputers such as ORNL's Summit to model aspects of the process. Those tools alone wouldn't be enough to capture the quantum nature of neutrinos.

"These neutrinos are entangled, which means they're interacting not just with their surroundings and not just with other neutrinos but with themselves," Savage said.

"It's extremely difficult to simulate this kind of system, because entanglement's an intrinsically quantum-mechanical property beyond what we can capture and approximate in classical computing. That's why we need a quantum computer that uses calculations based on [quantum physics](#) to model what's happening."

Savage and his co-author Marc Illa of the University of Washington's InQubator for Quantum Simulation obtained an allocation of time on Quantinuum's H1-1 quantum computer via QCUP, part of the Oak Ridge Leadership Computing Facility, which awards time on privately owned quantum processors around the country to support research projects. The Quantinuum computer uses trapped ions as qubits, one of several quantum computing approaches.

Classical computers store information in bits equal to either 0 or 1. In other words, a classical bit, like a light switch, exists in one of two states: on or off.

Quantum computers store information in qubits, the quantum equivalent of bits. Qubits, unlike classical bits, can exist in more than one state

simultaneously via quantum superposition—more like a dial with a wider range of more detailed settings than an on/off switch. That difference enables qubits to carry more information than classical bits. Scientists hope to use this increased capacity to fuel a quantum computing revolution built on a new generation of devices.

That capacity allowed Savage and the research team to simulate an approximation of the quantum-mechanical interactions between a supernova's neutrinos. An actual supernova would involve a minimum of a septendecillion, or  $10^{54}$ , neutrinos. Savage and Illa began their simulation using a simpler model with a system of 12 neutrinos.

Each neutrino "flavor," or type, found in nature corresponds to a "partner" particle: an electron, muon or tau. The model used in the study focused on just two flavors.

Quantum circuits—the quantum equivalent of traditional digital circuits—allowed the team to model the complicated connections and interactions between the particles so that each neutrino could interact with each of the others, not just its nearest neighbors.

The results offered a realistic approximation of how neutrinos become entangled at the quantum level, so that changing the properties of one also changes the properties of another. During a supernova, neutrinos can change flavor from an electron flavor to a muon flavor or to a tau flavor as the neutrinos begin to interact with each other and their surroundings. The detail provided by the simulations enabled the team to measure the evolution from one flavor to another over time of various entangled neutrinos.

Why track the flavor conversion? Because the mu and tau flavors of neutrinos interact differently with matter than their electron-flavored brethren. These interactions can impact the amounts and types of heavier

elements produced in the supernova explosion.

"These circuits turned out to approximate the neutrinos' behavior very well," Savage said. "We discovered we could use these simulations to measure neutrino entanglement in a statistically significant way and that we could identify a significant scaling in size as the number of neutrinos increased. This was the first time this kind of study had been done."

The primary hurdle for useful [quantum simulations](#) has been the relatively high error rate caused by noise that degrades qubit quality. The problem's so common the current generation of quantum computers has become known as noisy intermediate-scale quantum, or NISQ.

Various programming methods can help reduce these errors, but Savage and Illa didn't need those methods to conduct their study thanks to the high quality of the Quantinuum computer's qubits and gates. The computer's 12-qubit circuits proved to be sufficient for almost 200 of the 2-qubit gates.

"We found the systematic errors on the quantum hardware were less than the statistical errors," Savage said. "We still have a long way to go to predict the behavior of large neutrino systems with precision, and we don't know whether the current generation of NISQ devices can take us there. But this technique should be portable to other types of quantum computers, and the results help us set protocols that can be used to simulate larger systems of neutrinos."

Next steps include simulating a system of as many as 50 neutrinos. Savage hopes to model such systems in a variety of environments.

"We want to understand the implications of different thermal states, of states in and out of equilibrium," he said. "We're excited to see what we can explore."

**More information:** Marc Illa et al, Multi-Neutrino Entanglement and Correlations in Dense Neutrino Systems, *Physical Review Letters* (2023).  
[DOI: 10.1103/PhysRevLett.130.221003](https://doi.org/10.1103/PhysRevLett.130.221003)

Provided by Oak Ridge National Laboratory

Citation: Untangling the entangled: Quantum study shines fresh light on how neutrinos fuel supernovae (2024, June 22) retrieved 14 July 2024 from  
<https://phys.org/news/2024-06-untangling-entangled-quantum-fresh-neutrinos.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--