

July 23 2024, by Rod Boyce

New automated system provides a way to detect elusive volcanic vibrations



(a) Station map of the Alaska Volcano Observatory seismic and infrasound network at Pavlof Volcano. All seismic stations have a colocated infrasound sensor. Note that station PV6 was destroyed during the 2016 eruption, and is shown for reference because it was operational during the 2007, 2013, and 2014 eruptions. The inset map shows the location of Pavlof Volcano along the Alaska Peninsula. (b) Helicopter-based view of a small ash burst from Pavlof Volcano's active vent on 26 August 2022. The vents for the 2021–2022 eruption were located on the upper southeastern flank of the volcano relative to the summit. Credit: Valerie Wasser, University of Alaska Fairbanks

A new automated system of monitoring and classifying persistent



vibrations at active volcanoes can eliminate the hours of manual effort needed to document them.

Graduate student researcher Darren Tan at the University of Alaska Fairbanks Geophysical Institute led development of the system, which is based on machine learning. Machine learning is a branch of artificial intelligence focused on building systems that learn from data, identify patterns and make decisions with minimal human intervention.

Details about Tan's automated system are <u>published</u> in the journal *Journal of Geophysical Research: Solid Earth.*

His system documents volcanic tremor, a continuous, rhythmic seismic signal that emanates from a volcano. It often indicates underground movement of magma or gas and occurs regularly at <u>active volcanoes</u>.

Knowledge of volcanic tremor can help in forecasting and detecting eruptions.

Unlike volcanic earthquakes, volcanic tremor is a sustained ground rumble that can last from a few seconds to a year or more. It is primarily identified in spectrograms because of its varying intensity and frequency.

"Volcanic tremor isn't typically detected or catalogued, because it tends to be quite subtle in the <u>seismic data</u>," Tan said. "It doesn't have the impulsive onset like an earthquake does."

Detecting tremor is currently a manual process at the Alaska Volcano Observatory, with which Tan is also affiliated. The observatory is a joint program of the Geophysical Institute, the Alaska Division of Geological and Geophysical Surveys and the U.S. Geological Survey. Part of the observatory is based at the Geophysical Institute.



The observatory's daily duty seismologist scans spectrograms at 32 volcano-monitoring networks across Alaska, looking for the slight indications of tremor in addition to the obvious seismic signals.

"The duty seismologists go in every day, and sometimes twice a day or more depending on the <u>volcanic activity</u>, to look at spectrograms," Tan said. "They look from volcano to volcano, hour to hour, and it takes a long time."

Alaska has 54 volcanoes classified as "historically active," meaning they have erupted in the past approximately 300 years. Of those, 32 have seismic monitoring networks.

Tan drew upon the diversity of tremor signals from the 2021-2022 eruption of Pavlof Volcano, on the Alaska Peninsula, to build an extensive dataset of labeled seismic and low-frequency acoustic spectrograms. Those spectrograms represent a variety of classifications, such as tremor type, explosions and earthquakes, that were then used to train a computer model for each data type.

The trained models can detect and classify volcanic tremor in near real time. Humans will still be involved in interpreting what the automation produces, however.

"To be able to place our focus on time periods of interest, that is key," Tan said. "I think that reinvents the way we can monitor long-duration eruptions, because things can get missed when a volcano is active for a year and a half or two years."

"This automated method of detecting tremor is also an important contribution to the forecasting and detection of eruptions," he said.

Tan said machine learning is a rapidly growing field with great



possibilities.

"It's like the Wild West of <u>machine learning</u> right now," he said. "Everyone is trying to dip their toes into this, but it is important to do so carefully."

UAF researchers among the seven co-authors of the journal paper include David Fee, Alaska Volcano Observatory coordinating scientist at the Geophysical Institute; Társilo Girona, Geophysical Institute research assistant professor; and research assistant professor Taryn Lopez, also of the Geophysical Institute.

Matthew Haney, Chris Waythomas and Aaron Wech at the USGS and former UAF postdoctoral researcher Alex Witsil, now at Applied Research Associates in North Carolina, are also co-authors.

More information: Darren Tan et al, Detection and Characterization of Seismic and Acoustic Signals at Pavlof Volcano, Alaska, Using Deep Learning, *Journal of Geophysical Research: Solid Earth* (2024). DOI: 10.1029/2024JB029194

Provided by University of Alaska Fairbanks

Citation: New automated system provides a way to detect elusive volcanic vibrations (2024, July 23) retrieved 23 July 2024 from https://phys.org/news/2024-07-automated-elusive-volcanic-vibrations.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.