


Bright prospects for engineering quantum light

July 31 2024, by James Badham



Concept illustration depicting a quantum defect emitting a single photon. Credit: Mark Turianksy

Computers benefit greatly from being connected to the internet, so we might ask: What good is a quantum computer without a quantum internet?

The secret to our modern internet is the ability for data to remain intact

while traveling over long distances, and the best way to achieve that is by using photons.

Photons are single units ("quanta") of light. Unlike other quantum particles, photons interact very weakly with their environment. That stability also makes them extremely appealing for carrying [quantum information](#) over long distances, a process that requires maintaining a delicate state of entanglement for an extended period of time. Such photons can be generated in a variety of ways.

One possible method involves using atomic-scale imperfections (quantum defects) in crystals to generate single photons in a well-defined quantum state.

Decades of optimization have resulted in [fiber-optic cables](#) that can transmit photons with extremely low loss. However, this low-loss transmission works only for light in a narrow range of wavelengths, known as the "telecom [wavelength](#) band."

Identifying quantum defects that produce [photons](#) at these wavelengths has proven difficult. Researchers at the UC Santa Barbara College of Engineering conducted research to understand why that is and describe their findings in "[Rational Design of Efficient Defect-Based Quantum Emitters](#)," published in the journal *APL Photonics*.

"Atoms are constantly vibrating, and those vibrations can drain energy from a light emitter," says UCSB materials professor Chris Van de Walle. "As a result, rather than emitting a photon, a defect might instead cause the atoms to vibrate, reducing the light-emission efficiency."

Van de Walle's group developed [theoretical models](#) to capture the role of atomic vibrations in the photon-emission process and studied the role of various defect properties in determining the degree of efficiency.

Their work explains why the efficiency of single-photon emission drastically decreases when the emission wavelength increases beyond the wavelengths of visible light (violet to red) to the infrared wavelengths in the telecom band. The model also allows the researchers to identify techniques for engineering emitters that are brighter and more efficient.

"Choosing the host material carefully, and conducting atomic-level engineering of the vibrational properties are two promising ways to overcome low efficiency," said Mark Turiansky, a postdoctoral researcher in the Van de Walle lab, a fellow at the NSF UC Santa Barbara Quantum Foundry, and the lead researcher on the project.

Another solution involves coupling to a photonic cavity, an approach that benefited from the expertise of two other Quantum Foundry affiliates: computer engineering professor Galan Moody and Kamyar Parto, a graduate student in the Moody lab.

The team hopes that their model and the insights it provides will prove useful in designing novel quantum emitters that will power the quantum networks of the future.

More information: Mark E. Turiansky et al, Rational design of efficient defect-based quantum emitters, *APL Photonics* (2024). [DOI: 10.1063/5.0203366](https://doi.org/10.1063/5.0203366)

Provided by University of California - Santa Barbara

Citation: Bright prospects for engineering quantum light (2024, July 31) retrieved 9 September 2024 from <https://phys.org/news/2024-07-bright-prospects-quantum.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.