

# Cosmic simulation reveals how black holes grow and evolve

July 2 2024, by Kimm Fesenmaier

---



This still from the simulation shows a supermassive black hole, or quasar, surrounded by a swirling disk of material called an accretion disk. Credit: Caltech/Phil Hopkins group

A team of astrophysicists led by Caltech has managed for the first time to simulate the journey of primordial gas dating from the early universe

to the stage at which it becomes swept up in a disk of material fueling a single supermassive black hole. The new computer simulation upends ideas about such disks that astronomers have held since the 1970s and paves the way for new discoveries about how black holes and galaxies grow and evolve.

"Our new [simulation](#) marks the culmination of several years of work from two large collaborations started here at Caltech," says Phil Hopkins, the Ira S. Bowen Professor of Theoretical Astrophysics.

The first collaboration, nicknamed FIRE (Feedback in Realistic Environments), has focused on the larger scales in the universe, studying questions such as how galaxies form and what happens when galaxies collide. The other, dubbed STARFORGE, was designed to examine much smaller scales, including how stars form in individual clouds of gas.

"But there was this big gap between the two," Hopkins explains. "Now, for the first time, we have bridged that gap."

To do that, the researchers had to build a simulation with a resolution that is more than 1,000 times greater than the previous best in the field.

To the team's surprise, as [reported](#) in *The Open Journal of Astrophysics*, the simulation revealed that magnetic fields play a much larger role than previously believed in forming and shaping the huge disks of material that swirl around and feed the supermassive [black holes](#).

"Our theories told us the disks should be flat like crepes," Hopkins says. "But we knew this wasn't right because astronomical observations reveal that the disks are actually fluffy—more like an angel cake. Our simulation helped us understand that magnetic fields are propping up the disk material, making it fluffier."

## **Visualizing the activity around supermassive black holes using 'super zoom-ins'**

In the new simulation, the researchers performed what they call a "super zoom-in" on a single supermassive black hole, a monstrous object that lies at the heart of many galaxies, including our own Milky Way. These ravenous, mysterious bodies contain anywhere from thousands to billions of times the mass of the sun, and thus exert a huge effect on anything that comes near.

Astronomers have known for decades that as gas and dust are pulled in by the tremendous gravity of these black holes, they are not immediately sucked in. Instead, the material first forms a rapidly swirling disk called an accretion disk. And as the material is just about to fall in, it radiates a huge amount of energy, shining with a brilliance unmatched by just about anything in the universe. But much is still not known about these active [supermassive black holes](#), called quasars, and how the disks that feed them form and behave.

While disks around supermassive black holes have been imaged previously—the Event Horizon Telescope imaged disks circling black holes at the heart of our own galaxy in 2022 and Messier 87 in 2019—these disks are much closer and more tame than the ones that churn around quasars.

To visualize what happens around these more active and distant black holes, astrophysicists turn to supercomputer simulations. They feed information about the physics at work in these galactic settings—everything from the basic equations that govern gravity to how to treat dark matter and stars—into thousands of computing processors that work in parallel.

This input includes many algorithms, or series of instructions, for the

computers to follow to recreate complicated phenomena. So, for example, the computers know that once gas becomes dense enough, a star forms. But the process is not that straightforward.

"If you just say gravity pulls everything down and then eventually the gas forms a star and stars just build up, you'll get everything wildly wrong," Hopkins explains.


After all, stars do many things that affect their surroundings. They shine radiation that can heat up or push surrounding gas. They blow winds like the [solar wind](#) created by our own sun, which can sweep up material. They explode as supernovae, sometimes launching material clear out of galaxies or changing the chemistry of their surroundings. So, the computers must know all the ins and outs of this "stellar feedback" as well, as it regulates how many stars a galaxy can actually form.

## **Building a simulation that spans multiple scales**

But at these larger scales, the set of physics that are most important to include and what approximations can be made differ from those at smaller scales. For example, on the galactic scale, the complicated details of how atoms and molecules behave are extremely important and must be built into any simulation. However, scientists agree that when simulations focus on the more immediate area around a black hole, molecular chemistry can be mostly ignored because the gas there is too hot for atoms and molecules to exist. Instead, what exists there is hot ionized plasma.

Creating a simulation that could cover all the relevant scales down to the level of a single accretion disk around a supermassive black hole was a huge computational challenge—one that also required a code that could handle all the physics.

"There were some codes that had the physics that you needed to do the small-scale part of the problem and some codes that had the physics that you needed to do the larger, cosmological part of the problem, but nothing that had both," Hopkins says.



An earlier still from the simulation shows a tangle of merging galaxies. Credit: Caltech/Phil Hopkins group

The Caltech-led team used a code they call GIZMO for both the large- and small-scale simulation projects. Importantly, they built the FIRE project so that all the physics they added to it could work with the STARFORGE project, and vice versa.

"We built it in a very modular way, so that you could flip on and off any of the pieces of physics that you wanted for a given problem, but they were all cross compatible," Hopkins says.

This allowed the scientists in the latest work to simulate a black hole that is about 10 million times the mass of our sun, beginning in the [early universe](#). The simulation then zooms in on that black hole at a moment when a giant stream of material is torn off a cloud of star-forming gas and begins to swirl around the supermassive black hole. The simulation can continue zooming in, resolving a finer area at each step as it follows the gas on its way toward the hole.

## Surprisingly fluffy, magnetic disks

"In our simulation, we see this accretion disk form around the black hole," Hopkins says. "We would have been very excited if we had just seen that [accretion disk](#), but what was very surprising was that the simulated disk doesn't look like what we've thought for decades it should look like."

In two seminal papers from the 1970s that described the accretion disks fueling supermassive black holes, scientists assumed that thermal pressure—the change in pressure caused by the changing temperature of the gas in the disks—played the dominant role in preventing such disks from collapsing under the tremendous gravity they experience close to the black hole. They acknowledged that magnetic fields might play a minor role in helping to shore up the disks.

In contrast, the new simulation found that the pressure from the magnetic fields of such disks was actually 10,000 times greater than the pressure from the heat of the gas.

"So, the disks are almost completely controlled by the magnetic fields,"

Hopkins says. "The magnetic fields serve many functions, one of which is to prop up the disks and make the material puffy."

This realization changes a host of predictions scientists can make about such accretion disks, such as their mass, how dense and thick they should be, how fast material should be able to move from them into a black hole, and even their geometry (such as whether the disks can be lopsided).

Looking forward, Hopkins hopes this new ability to bridge the gap in scales for cosmological simulations will open many new avenues of research. For example, what happens in detail when two galaxies merge? What types of stars form in the dense regions of galaxies where conditions are unlike those in our sun's neighborhood? What might the first generation of stars in the universe have looked like?

"There's just so much to do," he says.

**More information:** Philip F. Hopkins et al, FORGE'd in FIRE: Resolving the End of Star Formation and Structure of AGN Accretion Disks from Cosmological Initial Conditions, *The Open Journal of Astrophysics* (2024). [DOI: 10.21105/astro.2309.13115](https://doi.org/10.21105/astro.2309.13115)

Provided by California Institute of Technology

Citation: Cosmic simulation reveals how black holes grow and evolve (2024, July 2) retrieved 2 July 2024 from <https://phys.org/news/2024-07-cosmic-simulation-reveals-black-holes.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.