

Grain boundaries weaken in planetary interiors, research suggests



Results indicate that grain boundary strength and motion, accommodated by either shear-coupled migration (SCM) or grain boundary sliding (GBS), strongly varies with pressure resulting in grain boundary hardening and weakening across a broad pressure range. Especially at pressure conditions of super-Earth exoplanets (~120–400 GPa), grain boundary weakening is observed with increasing depth during shear-coupled migration. Credit: S. Ritterbex and T. Tsuchiya

July 17 2024



Mantle convection and associated plate tectonics of planets like the Earth are governed by the deformation of mantle rocks. This deformation occurs through the motion of defects in the crystal lattices of minerals. Thus, the physical properties of these structural defects under pressure have profound implications on the dynamics of Earthlike planets.

A collaborative team of researchers, led by Dr. Sebastian Ritterex, a former post-doc of the Geodynamics Research Center, Ehime University and now a researcher of the Department of Earth Sciences, Utrecht University, applied massive parallel high-performance computer simulations based on quantum mechanical atomic-scale modeling to shed new light on the enigmatic behavior of grain boundaries under the extreme pressures that prevail in planetary interiors.

This theoretical methodology, called "ab initio simulations," enables the researchers to compute chemical bonding very accurately. It is a powerful tool for determining material properties under <u>extreme</u> <u>conditions</u> in the interior of planets where it is difficult to conduct experiments.

Based on the above theoretical mineral physics approach, the team examined the <u>mechanical behavior</u> and thermodynamic properties of high-angle tilt grain boundaries in (Mg,Fe)O ferropericlase, the second most abundant mineral in the Earth's lower <u>mantle</u> and possibly in the mantles of super-Earth exoplanets.

In this study, in addition to the standard density functional theory, the internally consistent LDA+U method was applied to reproduce the electronic structure of iron more accurately.

The results of the mechanical behavior indicate that the very high <u>pressure</u> conditions in terrestrial planets have a strong effect on the



mechanisms of grain boundary motion which govern intercrystalline deformation. The research proved for the first time that structural transformations of grain interfaces, induced by pressure with increasing depth in planetary mantles, trigger a change in the mechanism and in the direction of grain boundary motion.

The team also demonstrated that significant mechanical weakening of grain boundaries can develop under multi-megabar pressures. This is counterintuitive because it is usually thought that with increasing pressure, atomic arrangements in materials become more closely packed, making them harder.

This phenomenon of grain boundary weakening is caused by a change in the transition state structure of grain boundaries during their motion under extremely high pressures. Analyses of their data <u>presented</u> in the *Journal of Geophysical Research: Solid Earth* published in April 2024 identify grain boundary weakening in ferropericlase as one of the potential mechanisms for viscosity reductions with increasing depth in the mantle of super-Earth exoplanets.

The team performed additional thermodynamic modeling of the iron partitioning behavior between bulk and grain boundaries. They determined that <u>grain size</u> is an important factor in controlling the grain boundary segregation of iron in polycrystalline ferropericlase in the hot and dense lower mantle.

It is well known that incorporation of substitutional Fe(II) in bulk MgO has a significant effect on its physical properties such as density and seismic wave velocities, since Fe(II) undergoes an electronic spin transition at high pressure in the Earth's interior. There had been no previous information about the spin states of Fe(II) within grain boundaries.



The modeling now shows that the electronic spin state of Fe(II) within ferropericlase tilt grain boundaries is controlled by structural grain boundary transformations at high pressure in the Earth's lower mantle. This mechanism influences the pressure conditions of the iron spin crossover in polycrystalline (Mg,Fe)O with micrometer or smaller grain sizes.

The findings indicate that the iron spin crossover pressure in ferropericlase may increase by several tens of GPa due to pressureinduced structural grain boundary transitions in dynamically active finegrained lower mantle regions compared to more thermodynamically stable regions in the lower mantle.

While the team is happy with these breakthroughs, more systematic data from theoretical modeling as well as from experiments and electron microscopy observations will be needed to achieve better insights into the collective effects of grain boundaries on the rheological and thermodynamic properties of polycrystalline ferropericlase at the appropriate pressures and temperatures in planetary mantles.

More information: Sebastian Ritterbex et al, Atomic-Scale Study of Intercrystalline (Mg,Fe)O in Planetary Mantles: Mechanics and Thermodynamics of Grain Boundaries Under Pressure, *Journal of Geophysical Research: Solid Earth* (2024). DOI: 10.1029/2023JB028375

Provided by Ehime University

Citation: Grain boundaries weaken in planetary interiors, research suggests (2024, July 17) retrieved 17 July 2024 from <u>https://phys.org/news/2024-07-grain-boundaries-weaken-planetary-interiors.html</u>



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.