


What makes high temperature superconductivity possible? Researchers get closer to a unified theory

July 30 2024



Researchers combined high magnetic fields with X-ray scattering to reveal the connection between superconducting vortices (black circles), charge density waves (red wiggles), and spin density waves (blue wiggles) in a cuprate superconductor. Credit: Stanford University

In copper-containing materials called cuprates, superconductivity competes with two properties called magnetic spin and electric charge density wave (CDW) order. These properties reveal different parts of the



electrons in the superconductor. Each electron possesses spin and charge.

In a regular metal, the spins cancel each other out and electrical charges are uniform across a material. However, the strong electron–electron interactions in <u>high-temperature superconductors</u> such as cuprates give rise to other possible states.

New research <u>published</u> in *Nature Communications* has examined materials where strong magnetic interaction causes some of the electron spins to order along stripes. This occurs when spin density waves (SDW) and CDWs lock together to form a stable long-range "stripe state" where the peaks and valleys of the two waves are aligned.

This state reinforces the stability of the SDW and CDW. This stripe state competes with and interrupts the superconducting phase. Now, however, researchers have found that short-range CDW can be compatible, rather than competitive, with superconductivity in cuprate materials. This finding runs counter to scientific conventional wisdom.

As expected, short-range CDW competes with long-range stripes to suppress superconductivity. Unexpectedly, CDW also coexists with and is enhanced by short-range superconductivity.

The research also identified the possibility that short-range charge order may enable the formation and motion of vortices in the superconducting phase. This means researchers may be able to stabilize superconductivity at higher temperatures and magnetic fields by controlling or enhancing short-range charge order.

The results also provide important insight into the development of a unified quantum description of superconductivity in the cuprates.

The research was performed using X-ray measurements in a previously



uncharted high <u>magnetic field</u> regime of the cuprate $La_{1.885}Sr_{0.115}CuO_4$.

The CDW order consists of two components marked by distinct magnetic field and temperature dependencies.

The sample spontaneously segregates into superconducting and, alternatively, spin–charge <u>stripe</u> ordered regions, clarifying how long-range spin order and bulk superconductivity coexist.

More interestingly, the static vortex state observed at low fields can be made fluid—a vortex liquid state—in high fields ranging from 12 to 24 Tesla.

At these fields, the long-range superconducting phase is suppressed by field-induced mobile vortices. Surprisingly, a sudden enhancement to the CDW intensity is commensurate with the vortex melting field—a field much smaller than the upper critical field that quenches superconductivity.

This research supports the phase-disordering scenario for the superconducting transition, motivating a unified quantum description of density waves and <u>superconductivity</u> in <u>cuprate superconductors</u>.

More information: J.-J. Wen et al, Enhanced charge density wave with mobile superconducting vortices in $La_{1.885}Sr_{0.115}CuO_4$, *Nature Communications* (2023). DOI: 10.1038/s41467-023-36203-x

Provided by US Department of Energy

Citation: What makes high temperature superconductivity possible? Researchers get closer to a unified theory (2024, July 30) retrieved 30 July 2024 from <u>https://phys.org/news/2024-07-high-</u>



temperature-superconductivity-closer-theory.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.