


Results suggests titanium-48's nuclear structure changes when observed at varying distances

July 19 2024



The nuclear structure of titanium-48 changes from a shell model structure to an α -cluster structure based on the distance from the center of the nucleus. Credit: Osaka Metropolitan University

The world around us is made up of particles invisible to the naked eye, but physicists continue to gain insights into this mysterious realm.

Findings [published](#) in *Physical Review C* by Osaka Metropolitan University researchers show that the nuclear structure of an atom likely changes depending on the distance the protons and neutrons are from the center of the nucleus.

OMU graduate student Maito Okada, Associate Professor Wataru Horiuchi, and Professor Naoyuki Itagaki from the Graduate School of Science compared calculations using [theoretical models](#) with existing [experimental data](#) to determine whether titanium-48, the most common isotope of titanium with 22 protons and 26 neutrons, has a shell model structure or an α -cluster (alpha cluster) structure.

While shell models are symmetric, α -cluster structures are thought to have an [alpha particle](#) at the outer region of the [nucleus](#), making an asymmetrical configuration. An α -particle is the same as helium with two protons and two neutrons. In alpha decay, this particle is emitted; for example, titanium-48 becomes calcium-44 if such decay occurs.

The OMU team calculated the collision effect of high-energy accelerated protons and α -particles on titanium-48. This was based on a theory of nuclear reactions in which the impact of [protons](#) on a nucleus reflects the structure near the surface of the target nucleus, while the collision of α -particles on a nucleus reflects the structure of the outer regions.

The results suggest that titanium-48 changes from a shell model structure to an α -cluster structure depending on the distance from the center of the nucleus.

"These results upend the conventional understanding of nuclear structure and can be expected to provide clues to the α -decay process that occurs in heavy nuclei, which has not been solved for nearly 100 years," Professor Horiuchi said, referring to the Gamow theory on nuclear

decay.

"In the future, we would like to extend the results obtained through this research to take on the challenge of solving issues related to heavier nuclei."

More information: M. Okada et al, Shell-cluster transition in ^{48}Ti , *Physical Review C* (2024). DOI: [10.1103/PhysRevC.109.054324](https://doi.org/10.1103/PhysRevC.109.054324). On *arXiv*: arxiv.org/html/2403.01685v1

Provided by Osaka Metropolitan University

Citation: Results suggests titanium-48's nuclear structure changes when observed at varying distances (2024, July 19) retrieved 19 July 2024 from <https://phys.org/news/2024-07-results-titanium-nuclear-varying-distances.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.