

Team develops a technique to detect nutrients in soil faster and more affordably

July 22 2024, by Janese Heavin



Researchers test a new method to more efficiently measure ions in soil. Credit: University of Missouri

Every year, Missouri farmers lose millions of dollars in valuable nutrients that wash away into rivers and lakes. These nutrients—nitrates



and phosphates found in fertilizers—are crucial for plant growth, but they wreak havoc on aquatic life when they end up in waterways.

University of Missouri researcher Matthias Young believes he's found a way to keep the nutrients on land where they belong. He's developing a technique that could lead to new types of sensors to help farmers better measure nitrates and phosphates in the soil, allowing them to use <u>fertilizers</u> more efficiently.

"Nitrates and phosphates released into waterways cause algae blooms that kill fish and hurt tourism in waterfront communities," said Young, an assistant professor of chemical engineering with a joint appointment in chemistry in Mizzou's College of Arts and Science. "Not only that, but farmers don't want nitrates and phosphates washing off their fields. Based on data from the Environmental Protection Agency and the United States Department of Agriculture, an estimated \$100 million of fertilizer is wasted every year in Missouri alone. This work helps fish, wildlife, tourism and farmers."

A new path forward

While his team is still working out some kinks, Young's research shows promise for a faster, more reliable and cost-effective way to measure ions such as nitrates in the soil.

That's important because if nitrate levels are low, plants won't get the nutrients they need to grow. But if they're too high, adding more fertilizer is wasteful because excess nitrates will run off into waterways.

Existing devices, known as ion-selective electrodes, can measure <u>nitrates</u> and <u>phosphates</u>, but they're costly and imprecise. One current method requires farmers to collect <u>soil samples</u> and send them to a lab for analysis, which is slow and only represents a small patch of land. Labs



have access to more sophisticated sensors that rely on special molecules to bind to and measure ions, but these molecules can run about \$10,000 per gram.

"Farmers are deciding how much fertilizer to deliver, but they're making decisions based on limited information," Young said. "Even the expensive sensors have to be recalibrated frequently, meaning they can't be left in the field."

Young's new design changes the way information is collected.

He and his team are devising a sensor that measures how quickly ions move through a membrane, rather than how strongly they bind to it. Using a pulsed electrical current, these devices are more sensitive and reliable.

Young is consulting with Missouri farmers as he makes progress in hopes of commercializing sensors that are inexpensive and practical.

"We're developing handheld sensors to collect measurements in realtime, as well as sensors that can be installed in the field," Young said. "They're not yet ready to deploy, as we're still having some challenges, but it is offering a new path forward."

He recently outlined his <u>findings</u> in *ACS Omega*. Co-authors were Ryan Gettler, Shima Mehregan, Henry Koeing and Andie Kaess.

More information: Ryan C. Gettler et al, Nonequilibrium Anion Detection in Solid-Contact Ion-Selective Electrodes, *ACS Omega* (2024). DOI: 10.1021/acsomega.4c00131



Provided by University of Missouri

Citation: Team develops a technique to detect nutrients in soil faster and more affordably (2024, July 22) retrieved 25 August 2024 from <u>https://phys.org/news/2024-07-team-technique-nutrients-soil-faster.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.