


Researchers identify unique phenomenon in Kagome metal

July 30 2024, by Amy Walden



A diagram of the Kagome metal cesium vanadium antimonide showing plasmon

waves moving through the material. Credit: Guangxin Ni

In traditional Japanese basket-weaving, the ancient "Kagome" design seen in many handcrafted creations is characterized by a symmetrical pattern of interlaced triangles with shared corners. In quantum physics, the Kagome name has been borrowed by scientists to describe a class of materials with an atomic structure closely resembling this distinctive lattice pattern.

Since the latest family of Kagome metals was discovered in 2019, physicists have been working to better understand their properties and potential applications. A new study led by Florida State University Assistant Professor of Physics Guangxin Ni focuses on how a particular Kagome metal interacts with light to generate what are known as [plasmon](#) polaritons—nanoscale-level linked waves of electrons and [electromagnetic fields](#) in a material, typically caused by light or other electromagnetic waves. The work was [published](#) in *Nature Communications*.

Previous research has examined plasmons in regular metals, but not as much in Kagome metals, where the behavior of electrons is more complex. In this study, the FSU researchers examined the metal cesium vanadium antimonide, also known by its chemical formula CsV_3Sb_5 , to better understand the properties that make it a promising contender for more precise and efficient photonic technologies.

The researchers identified for the first time the existence of plasmons in CsV_3Sb_5 and found that the wavelength of those plasmons depends upon the thickness of the metal.

They also found that changing the frequency of a laser shining at the

metal caused the plasmons to behave differently, turning them into a form known as "hyperbolic bulk plasmons," which spread through the material rather than staying confined to the surface. As a result, these waves lost less energy than before, meaning they could travel more effectively.

"Hyperbolic plasmon polaritons are rare in natural metals, but our research reveals how electron interactions can create these unique waves at the nanoscale," Ni said. "This breakthrough is key for advancing technologies in nano-optics and nano-photonics."

To explore how plasmons interacted with the metal, the researchers grew single crystals of CsV_3Sb_5 and then placed thin flakes of the material onto specially prepared gold surfaces. By using lasers to perform scanning infrared nano-imaging, they observed how the metal's plasmon polaritons—waves of electrons interacting with electromagnetic fields—changed in interesting ways.

"What makes CsV_3Sb_5 interesting is how it interacts with light on a very small scale, what's known as nano-optics," said lead author Hossein Shiravi, a graduate research assistant at the FSU-headquartered National High Magnetic Field Laboratory. "We found that over a wide range of infrared light frequencies, the correlated electrical properties within the metal triggered the formation of hyperbolic bulk plasmons."

That hyperbolic pattern means less energy is lost. The team's findings reveal new information about the way Kagome metal CsV_3Sb_5 behaves under various conditions, providing researchers with a more accurate picture of its properties and potential real-world applications.

"Hyperbolic plasmon polaritons can offer a range of amazing nano-optical features and abilities," Ni said. "They have the potential to boost optical communication systems, allow for super-clear imaging beyond

current limits and make photonic devices work better. They could also be useful for sensing things like environmental changes and medical diagnostics because they react strongly to their surroundings. These qualities make them key for advancing future optical and photonic technologies."

The CsV_3Sb_5 metal was a promising choice for plasmon research because of its unusual electronic and optical properties, such as its potential ability to force waves of plasmons to move in a single direction, to name just one. Recent advances in imaging technology at the nano-scale level helped the researchers complete their work.

"Electronic losses typically encountered in conventional metals have previously complicated efforts to observe exotic light-matter coupling effects, including hyperbolic polaritons," Ni said. "This is part of what makes this an exciting breakthrough. It will be interesting to continue exploring nano-optical phenomena in unconventional metals owing to their potential to contribute to future technologies."

FSU graduate student Aakash Gupta was also a co-author on this study. The study was conducted in collaboration with researchers from the University of California Santa Barbara, Oak Ridge National Laboratory in Tennessee, Tsinghua University in China, and Germany's University of Stuttgart, Leipzig University, and Institute of Ion Beam Physics and Materials Research.

More information: H. Shiravi et al, Plasmons in the Kagome metal CsV_3Sb_5 , *Nature Communications* (2024). [DOI: 10.1038/s41467-024-49723-x](https://doi.org/10.1038/s41467-024-49723-x)

Provided by Florida State University

Citation: Researchers identify unique phenomenon in Kagome metal (2024, July 30) retrieved 9 September 2024 from <https://phys.org/news/2024-07-unique-phenomenon-kagome-metal.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.