


Cultured zooplankton provide new avenue to monitor and rid oceans of microplastic pollutants

August 28 2024



Graphical abstract. Credit: *Science of The Total Environment* (2024). DOI: 10.1016/j.scitotenv.2024.174279

Testing for levels of microplastic contamination in marine plankton aims

to help develop new ways to assess and manage the rising global pollution problem, Flinders University experts say.

Researchers examined the effects of five different chemical digestive aids on common plastics using low, medium and high levels of zooplankton and have published the results in an article in [*Science of the Total Environment*](#).

Plastic pollution is the leading form of marine litter, with ever-increasing levels of plastic waste pouring into streams and oceans every year, says Flinders University Ph.D. candidate Elise Tuuri.

"Plastic pollution is found in deep-sea and shoreline sediments, [surface waters](#) and in marine biota. This has far-reaching impacts on ecosystems, including harm to [marine animals](#), and has the potential to impact [food webs](#) and habitats," she says.

"If scientists can find a way to determine the association of microplastics with zooplankton, we have a chance to understand and potentially curb damage from microplastics."

Microplastics (particles less than 5mm) are turning up in the guts of fish and shellfish, and there is concern about the safety of seafood. They can also occur in drinking water and other food products. Both plastics and chemical additives can be toxic.

The exponential growth of plastic production from 2 million metric tons in 1950 to 380m metric tons in 2015 has made [plastic pollution](#) the predominant form of anthropogenic marine litter globally. Production is set to triple by 2050.

Using cultured zooplankton kept in controlled conditions, the Flinders University researchers were able to examine the effects of five different

chemical digestive aids—acid, two different alkaline, enzymatic and oxidative—on common plastics (polyamide, polyethylene, polyethylene terephthalate, polypropylene, and polystyrene).

Professor Sophie Leterme, co-author of the study says the various methods resulted in varying degrees of damage to the chemical and physical makeup of these commonly found environmental microplastic pollutants.

Leterme, director of the ARC Center for Biofilm Research and Innovation, says reporting microplastic abundances via these methods could help with the collection of data about [microplastic](#) particles.

"This will enable us to understand the environmental and [health impacts](#), and develop effective strategies to mitigate marine pollution," she says.

More information: Elise M. Tuuri et al, Efficacy of chemical digestion methods to reveal undamaged microplastics from planktonic samples, *Science of The Total Environment* (2024). [DOI: 10.1016/j.scitotenv.2024.174279](#)

Provided by Flinders University

Citation: Cultured zooplankton provide new avenue to monitor and rid oceans of microplastic pollutants (2024, August 28) retrieved 28 August 2024 from <https://phys.org/news/2024-08-cultured-zooplankton-avenue-oceans-microplastic.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--