

'Mirror' nuclei help connect nuclear theory and neutron stars

August 13 2024



Charge radii differences in mirror nuclei, which have opposite numbers of protons and neutrons, can help constrain parameters for the equation of state nuclear matter, which describes the properties of astrophysical objects such as neutron stars. Credit: Ronald Garcia/Dean Lee

Adding or removing neutrons from an atomic nucleus leads to changes in



the size of the nucleus. This in turn causes tiny changes in the energy levels of the atom's electrons, known as isotope shifts. Scientists can use precision measurements of these energy shifts to measure the radius of the nucleus of an isotope.

In a recent study, researchers made laser-assisted measurements of the nuclear radii of the stable silicon isotopes silicon-28, silicon-29, and silicon-30. They also measured the radius of the unstable silicon-32 nucleus, which has 14 protons and 18 <u>neutrons</u>. The researchers used the difference between the radius of the silicon-32 nucleus and its mirror nucleus, argon-32, which has 18 protons and 14 neutrons, to set limits on variables that help to describe the physics of astrophysical objects such as <u>neutron stars</u>. The results are an important step in developing nuclear theory, the study of nuclei and their components.

The paper is **<u>published</u>** in the journal *Physical Review Letters*.

Despite progress in nuclear theory, scientists still face long-standing challenges in their understanding of nuclei. For instance, researchers have not connected the description of nuclear size with the underlying theory of the strong nuclear force. Moreover, it is not clear whether nuclear theories that describe finite atomic nuclei can provide a reliable description of nuclear matter. This special form of matter consists of interacting protons and neutrons. Nuclear matter includes matter in extreme conditions such as neutron stars. Precision measurements of charge radii—the radius of atomic nuclei—help solve these open questions.

The researchers used laser spectroscopy measurements of atomic isotope shifts to measure the nuclear radius of different silicon isotopes at the BEam COoler and LAser spectroscopy facility (BECOLA) at the Facility for Rare Isotope Beams (FRIB) at Michigan State University.



The results provide an important benchmark for the development of nuclear theory. The charge radii difference between the silicon-32 nucleus and its mirror nucleus argon-32 was used to constrain parameters needed to describe the properties of dense neutron matter within neutron stars. The obtained results agree with the constraints from gravitational wave observations and other complementary observables.

More information: Kristian König et al, Nuclear Charge Radii of Silicon Isotopes, *Physical Review Letters* (2024). DOI: <u>10.1103/PhysRevLett.132.162502</u>. On *arXiv*: DOI: <u>10.48550/arxiv.2309.02037</u>

Provided by US Department of Energy

Citation: 'Mirror' nuclei help connect nuclear theory and neutron stars (2024, August 13) retrieved 13 August 2024 from <u>https://phys.org/news/2024-08-mirror-nuclei-nuclear-theory-neutron.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.