

# Plant signaling pathways decoded using 'optogenetic' tobacco plants

August 28 2024



Transgenic tobacco leaves after 24 hours of green light exposure: a, plant with anion channel GtACR1; green light-induced membrane potential change (depolarization) is shown on the left. b, plant with calcium ion channel XXM 2.0; green light-induced depolarization is shown on the right. Horizontal bar indicates exposure time. Credit: Meiqi Ding / JMU

Using newly generated "optogenetic" tobacco plants, research teams from the University of Würzburg's Departments of Plant Physiology and Neurophysiology have investigated how plants process external signals.

When it comes to survival, plants have a huge disadvantage compared to

many other living organisms: they cannot simply change their location if predators or pathogens attack them or the environmental conditions change to their disadvantage.

For this reason, plants have developed different strategies with which they react to such attacks. Such reactions are usually triggered by certain signals from the environment. As has long been known, the intracellular calcium concentration plays an important role in the processing of these signals.

However, in addition to changes in the cytoplasmic calcium level, changes in the cell's membrane potential have also been suspected as a signal transmitter. Research groups from the Departments of Neurophysiology, Pharmaceutical Biology and Botany at Julius-Maximilians-Universität Würzburg (JMU) have investigated the calcium-membrane potential relationship in more detail. They have now [published](#) their findings in the journal *Nature*.

## **Light-sensitive channels enable targeted manipulations**

For their study, the research teams worked with tobacco plants that carry [ion channels](#) that can be specifically switched on with light. More than 20 years ago, Peter Hegemann, Georg Nagel and Ernst Bamberg initiated the success of optogenetics, with their discovery and characterization of light-activated ion channels, so-called channelrhodopsins.

With the help of these light-sensitive proteins, which are obtained from algae and microorganisms, the JMU researchers were able to experimentally investigate whether the influx of calcium ions or anion efflux-mediated depolarization of the cell membrane is decisive for the plant's reaction to a certain stress situation. However, the scientists had

to do a great deal of preparatory work before they were able to do this.

## **Optogenetics with rhodopsins**

Channelrhodopsins, ion channels that carry an intrinsic rhodopsin-based light switch, revolutionized neuroscience through the light-controlled investigation of neuronal networks. The use of channelrhodopsins in plant research only became possible 20 years later, through close collaboration between the group of Georg Nagel, Professor at the Institute of Physiology at JMU, and plant researchers from the Würzburg Chairs of Botany 1, 2 and Pharmaceutical Biology.

In 2021, Georg Nagel's group, together with Dr. Kai Konrad, group leader at the JMU Chair of Prof. Rainer Hedrich Botany 1, published an approach to optimize the use of channelrhodopsins in plants by overcoming three main difficulties.

"Like all rhodopsins, including those in our eyes, channelrhodopsins require the small molecule retinal, also known as vitamin A, to absorb light. We humans get retinal mainly from beta-carotene, the provitamin A. However, [land plants](#) do not contain retinal, but a lot of beta-carotene," explains Dr. Shiqiang Gao, co-author of the publication and 'rhodopsin engineer' from the Optogenetics lab of the Department of Neurophysiology at JMU.

In 2021, Gao succeeded for the first time in combining the expression of channelrhodopsins with the production of retinal from beta-carotene in plant cells. This enabled the development of tobacco plants with a high retinal content and successful expression of channelrhodopsins.

Dr. Markus Krischke from the Metabolomics Core Unit at the Department of Pharmaceutical Biology headed by Professor Martin Müller at JMU Würzburg confirmed the high retinal content of the

various transgenic tobacco plants.

Comparable transgenic tobacco plants were produced for the recently published study by Dr. Meiqi Ding from the Department of Botany I under the direction of plant physiologist and expert for plant signal processing Dr. Konrad from the group of Professor Hedrich at the Department of Botany I.

"Most rhodopsins are activated by blue or green light. However, this is always a component of white light," explains Georg Nagel. As a result, the tobacco plants could not be grown in a greenhouse or under artificial white light, as is usually the case. Only in special growth chambers with red LED light, which can be used photosynthetically, it was possible to avoid unwanted rhodopsin activation.

Tests under different growth conditions showed: "Tobacco develops healthily and unchanged under red light compared to greenhouse conditions," says Dr. Konrad.

The expression of channelrhodopsin in tobacco cells often causes difficulties. In 2021, the Würzburg team of scientists succeeded in expressing the light-activated anion channel GtACR1 in tobacco plant cells. As a result, Georg Nagel's team was able to develop various channelrhodopsins that were optimized for the permeability of calcium ions. Finally, Dr. Gao and Dr. Shang Yang, both members of Nagel's group, succeeded in developing a very good calcium-conducting channelrhodopsin XXM 2.0 for targeted expression in tobacco plants.

This was the breakthrough: "The successful expression of channelrhodopsins with different ion selectivity in plant cells enables the comparison of different ion signals in parallel to the electrical signal, the so-called depolarization," explains Dr. Meiqi Ding. She used the calcium-conducting channelrhodopsin XXM 2.0 and the light-activated anion

channel GtACR1 to investigate the different ion signaling pathways in tobacco.

## **A new era in plant research**

These newly generated "optogenetic" [tobacco plants](#) made it possible to clarify the question of whether calcium influx or membrane depolarization is decisive for the plant's response to a specific stress situation.

"The answer was clear," says Dr. Konrad, corresponding author.

First author Dr. Ding from Dr. Konrad's group explains, "After activation of the anion channel, the leaves wilted and responded with the typical plant response to drought; the plant hormone abscisic acid (ABA) was produced and gene expression was ramped up to protect against desiccation."

"However, in the plants with the calcium channel, there was no change in ABA levels after optogenetic stimulation," Dr. Ding continued.

"Instead, the plants produced signal molecules and plant hormones to initiate defense mechanisms against predators, recognizable by white spots on the leaves," said Dr. Konrad.

Dr. Sönke Scherzer at the chair of Prof Hedrich was able to show by direct ROS measurements that reactive oxygen species (ROS) are released in the process.

Dirk Becker and Rainer Hedrich at the Chair of Botany 1, designed an experimental approach to support the working hypothesis using transcriptomic and bioinformatic analysis.

The scientists are convinced that this study is just the beginning of a new era in plant research. Ultimately, the signaling pathways of plants can now be better "illuminated" using various rhodopsins.

**More information:** Meiqi Ding et al, Probing plant signal processing optogenetically by two channelrhodopsins, *Nature* (2024). [DOI: 10.1038/s41586-024-07884-1](https://doi.org/10.1038/s41586-024-07884-1).  
[www.nature.com/articles/s41586-024-07884-1](https://www.nature.com/articles/s41586-024-07884-1)

Provided by Julius-Maximilians-Universität Würzburg

Citation: Plant signaling pathways decoded using 'optogenetic' tobacco plants (2024, August 28) retrieved 28 August 2024 from <https://phys.org/news/2024-08-pathways-decoded-optogenetic-tobacco.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--