


Scientists propose gliding arc discharge plasma seawater sterilization technology for mariculture

August 19 2024



The overview of GADP setup. (a) The schematic diagram of the experimental setup; (b) The usage scenario of GADP during the experiment and container diagram for three ASW volumes; (c) The structure diagram and cutaway view of the reactor chamber; (d) The 3D structure diagram and discharge principle of the electrodes in the reactor chamber. Credit: SIBET

Atmospheric pressure plasma technology has demonstrated vast potential in microbial inactivation due to its unique advantages, including no chemical residue, environmental friendliness, high treatment efficiency, low energy consumption and ease of use.

Researchers at the Suzhou Institute of Biomedical Engineering and Technology (SIBET) of the Chinese Academy of Sciences have proposed a strategy to sterilize mariculture utilizing gliding arc discharge plasma (GADP).


They independently developed an air GADP device that can work in seawater with different concentrations of *Vibrio parahaemolyticus* (VP).

VP is the most prevalent pathogen in intensive mariculture worldwide, posing a significant threat to both the aquaculture industry and public health safety. Currently, antibiotics are the primary means of prevention and control of VP infections in mariculture. However, extensive and [overuse of antibiotics](#) inevitably leads to issues such as antibiotic residues in seafood and bacterial resistance.


Optical emission spectroscopy (OES) diagnostics indicate that the GADP is rich in reactive species such as hydroxyl radicals, excited state oxygen atoms, [singlet oxygen](#), and nitric oxide.

In their new study, the SIBET researchers conducted a dose-effect study on the inactivation of VP in varying volumes of artificial seawater using GADP. Results showed that the GADP treatment achieved complete inactivation of VP in a 5000 mL volume within 20 minutes.

Results of the study titled "The effectiveness of gliding arc discharge plasma in sterilizing artificial seawater contaminated with *Vibrio parahaemolyticus*" have been [published](#) in the *Journal of Hazardous Materials*.



The inactivation effect of GADP against VP in 50 mL, 500 mL and 5000 mL of ASW, the concentrations of VP (a) 1.0×10^3 CFU/mL; (b) 5.0×10^3 CFU/mL; (c) 10×10^3 CFU/mL. Credit: SIBET



The potential sterilization mechanisms of GADP in artificial seawater. Credit: SIBET

Moreover, the data of the pH, salinity, temperature, and nitrate content measurements before and after treatment indicate that the GADP has no significant impact on the key water quality parameters of artificial seawater.

The researchers studied the potential sterilization mechanisms of GADP in artificial seawater by examining [hydroxyl radicals](#), [hydrogen peroxide](#), oxidation-reduction potential, intracellular reactive oxygen species, and cell membrane permeability.

This work provides a viable solution for infection with the halophilic pathogen *Vibrio parahaemolyticus* and demonstrates the potential of GADP in [seawater](#) sterilization.

More information: Meng-Ru Du et al, The effectiveness of gliding arc discharge plasma in sterilizing artificial seawater contaminated with *Vibrio parahaemolyticus*, *Journal of Hazardous Materials* (2024). [DOI: 10.1016/j.jhazmat.2024.135015](https://doi.org/10.1016/j.jhazmat.2024.135015)

Provided by Chinese Academy of Sciences

Citation: Scientists propose gliding arc discharge plasma seawater sterilization technology for mariculture (2024, August 19) retrieved 20 August 2024 from <https://phys.org/news/2024-08-scientists-gliding-arc-discharge-plasma.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.