


Scientists show the sun's magnetic activity influences the determination of its seismic age

August 19 2024



Evolution of the sun during cycle 23. The image shows a collage of 11 photographs of the sun taken by the SOHO satellite and highlights the 11-year solar activity cycle with minima around 1996 and 2006 and a maximum around 2001. Credit: SOHO/ESA/NASA

An international team of astronomers, led by a researcher from the University of Geneva (UNIGE), has shown that the sun's magnetic activity has a significant influence on its seismic characterization, contrary to predictions in previous studies. Important data such as its size, age and chemical composition depend on it.

These results pave the way for in-depth research to better understand the nature of the magnetic activity and its impact on stellar oscillations. The study is [published](#) in the journal *Astronomy & Astrophysics*.

Asteroseismology, or helioseismology in the case of the sun, is a fascinating branch of astronomy that studies the oscillations of stars.

"To understand it, you must imagine a star as a big ball of gas in constant motion. Inside this star, there are waves or pulsations that make it vibrate, rather like the sound that resonates in a musical instrument," explains Jérôme Bétrisey, a postdoctoral researcher in the Astronomy Department of the Faculty of Science at the UNIGE and first author of the study.

"These vibrations cause the star's surface to move slightly and change luminosity regularly. Thanks to very precise instruments, we can detect these variations in luminosity from Earth or from space," continues the researcher.

Listening to the symphony of the stars

By observing these changes, researchers can learn a great deal about the star's internal structure and determine important characteristics such as its size, age, chemical composition or stage in the stellar life [cycle](#). A precise understanding of the characteristics of stars is essential, among

other things, to determine the properties of the planets that surround them or to retrace the history of the Milky Way.

Despite some major successes in recent decades, asteroseismology has also shown that there are significant differences between observations and the predictions of theoretical models of the inner workings of stars.

Over the years, various methods have been developed to reduce these discrepancies, with varying degrees of success. However, none of the current methods consider the magnetic activity of stars, its impact on the results being supposed to be negligible.

The international team led by Bétrisey has just demonstrated the opposite. It has established that the age of the sun, as determined by helioseismology, varies significantly according to the level of activity of the solar cycle.

To give an order of magnitude, the sun is around 4.6 billion years old, and variations of up to 300 million years have been observed between solar minima. Although these variations may seem small compared with the age of the sun, they are no longer negligible given the level of precision that will be achieved by [future space missions](#).

Seismic age of sun correlates with the solar activity cycle

To better understand the impact of magnetic activity on the sun, the scientists analyzed 26.5 years of solar data, covering two complete activity cycles. They divided this data into about 90 small one-year series, each three months apart. For each of these series, a seismic analysis was carried out, making it possible to measure the evolution of the sun's fundamental properties such as its mass, radius and age over

time.

Two independent data sets were used to check the robustness of the results. One came from the University of Birmingham's BiSON (Birmingham Solar Oscillations Network) network of ground-based telescopes, and the other from the GOLF (Global Oscillations at Low Frequencies) instrument on board the SOHO (Solar and Heliospheric Observatory) satellite, which has been orbiting the sun since the mid-1990s.

Regardless of the configuration tested, the age of the sun determined by helioseismology was correlated with the level of activity of the solar cycle. They measured variations of around 6% on average between the periods of solar minimum and maximum, which is very significant in terms of the precision targeted for future space missions that will analyze other similar stars.

For example, the PLATO (PLAnetary Transits and Oscillations of stars) mission is aiming for a precision of 10% for the age of a star like the sun.

The study of the GOLF and BiSON data also showed that the impact of the activity cycle on the seismic age is more marked for the more active of the two cycles studied. This result is logical and expected from a physical point of view.

"However, the sun is not a particularly active star, which suggests that the impact of magnetic activity could be very significant for more active stars such as those that PLATO will detect," adds Bétrisey.

A bright future for the study of the magnetic activity of stars

The results of this study show that the magnetic activity of stars poses a major challenge for future space missions such as PLATO, particularly in terms of characterizing the most active stars. "However, this discovery also opens many exciting research prospects," concludes Bétrisey.

The magnetic activity of stars has a significant influence on [stellar oscillations](#), which complicates the precise determination of fundamental properties such as the mass, radius and age of stars. For future space missions, this means that it will be necessary to develop more sophisticated methods to take account of this magnetic impact.

The challenges posed by magnetic activity will also encourage researchers to explore this phenomenon further. This could lead to a better understanding of stellar physics, particularly in terms of how magnetic fields interact with the internal oscillations of stars. This research could also improve our understanding of stellar activity cycles, which are like solar cycles.

More information: J. Bétrisey et al, Imprint of the magnetic activity cycle on solar asteroseismic characterisation based on 26 years of GOLF and BiSON data, *Astronomy & Astrophysics* (2024). [DOI: 10.1051/0004-6361/202451365](#)

Provided by University of Geneva

Citation: Scientists show the sun's magnetic activity influences the determination of its seismic age (2024, August 19) retrieved 19 August 2024 from <https://phys.org/news/2024-08-scientists-sun-magnetic-seismic-age.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.