


New view of North Star reveals spotted surface

August 20 2024



CHARA Array false-color image of Polaris from April 2021 that reveals large

bright and dark spots on the surface. Polaris appears about 600,000 times smaller than the full moon in the sky. Credit: Georgia State University / CHARA Array

Researchers using Georgia State University's Center for High Angular Resolution Astronomy (CHARA) Array have identified new details about the size and appearance of the North Star, also known as Polaris. The new [research](#) is published in *The Astrophysical Journal*.


Earth's North Pole points to a direction in space marked by the North Star. Polaris is both a navigation aid and a remarkable star in its own right. It is the brightest member of a triple-star system and is a pulsating variable star. Polaris gets brighter and fainter periodically as the star's diameter grows and shrinks over a four-day cycle.

Polaris is a kind of star known as a Cepheid variable. Astronomers use these stars as "standard candles" because their true brightness depends on their period of pulsation: Brighter stars pulsate slower than fainter stars. How bright a star appears in the sky depends on the star's true brightness and the distance to the star. Because we know the true brightness of a Cepheid based on its pulsational period, astronomers can use them to measure the distances to their host galaxies and to infer the expansion rate of the universe.

A team of astronomers led by Nancy Evans at the Center for Astrophysics | Harvard & Smithsonian observed Polaris using the CHARA optical interferometric [array](#) of six telescopes at Mount Wilson, Calif. The goal of the investigation was to map the orbit of the close, faint companion that orbits Polaris every 30 years.

"The small separation and large contrast in brightness between the two stars makes it extremely challenging to resolve the binary system during

their [closest approach](#)," Evans said.



The CHARA Array is located at the Mount Wilson Observatory in the San Gabriel Mountains of southern California. The six telescopes of the CHARA Array are arranged along three arms. The light from each telescope is transported through vacuum pipes to the central beam combining lab. All the beams converge on the MIRC-X camera in the lab. Credit: Georgia State University

The CHARA Array combines the light of six telescopes that are spread across the mountaintop at the historic Mount Wilson Observatory. By combining the light, the CHARA Array acted like a 330-meter telescope to detect the faint companion as it passed close to Polaris. The observations of Polaris were recorded using the MIRC-X camera built by astronomers at the University of Michigan and Exeter University in the U.K. The MIRC-X camera has the remarkable ability to capture details of stellar surfaces.

The team successfully tracked the orbit of the close companion and measured changes in the size of the Cepheid as it pulsated. The [orbital motion](#) showed that Polaris has a mass five times larger than that of the sun. The images of Polaris showed that it has a diameter 46 times the size of the sun.

The biggest surprise was the appearance of Polaris in close-up images. The CHARA observations provided the first glimpse of what the surface of a Cepheid variable looks like.

"The CHARA images revealed large bright and [dark spots](#) on the surface of Polaris that changed over time," said Gail Schaefer, director of the CHARA Array. The presence of spots and the rotation of the star might be linked to a 120-day variation in measured velocity.

"We plan to continue imaging Polaris in the future," said John Monnier, an astronomy professor at the University of Michigan. "We hope to better understand the mechanism that generates the spots on the surface of Polaris."

The new observations of Polaris were made and recorded as part of the open access program at the CHARA Array, where astronomers from around the world can apply for time through the National Optical-Infrared Astronomy Research Laboratory (NOIRLab).

More information: Nancy Remage Evans et al, The Orbit and Dynamical Mass of Polaris: Observations with the CHARA Array, *The Astrophysical Journal* (2024). [DOI: 10.3847/1538-4357/ad5e7a](https://doi.org/10.3847/1538-4357/ad5e7a)

Provided by Georgia State University

Citation: New view of North Star reveals spotted surface (2024, August 20) retrieved 20 August 2024 from <https://phys.org/news/2024-08-view-north-star-reveals-surface.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--