


# Modeling study explains why amazon is such a biodiverse paradise

September 11 2024



Biomes reconstruction during the LGM (left) as modeled in Sato et al10. (middle) using bias correction and (right) using clustering, driven by an ensemble of LGM climate reconstructions. Credit: *npj Biodiversity* (2024). DOI:

10.1038/s44185-024-00056-4

The Amazon rainforest is home to a remarkable variety of plants and animals not found anywhere else on Earth, with some species only located in certain areas, but the reason for this has perplexed and divided scientists for decades.

Now a new international study, led by the UK Centre for Ecology & Hydrology (UKCEH), challenges traditional thinking about how the Amazon evolved during the last Ice Age, which spanned the period between around 2.6 million and 11,700 years ago.

It demonstrates that the world's largest tropical rainforest is more sensitive to environmental change than previously thought, providing a further warning about how the ongoing, large-scale, rapid human-driven climate and [land use change](#) presents a threat to this precious ecosystem.

Some scientists have argued that, during the last Ice Age, the Amazon had "forest islands" (referred to as "refugia") that were completely isolated, while others have suggested the forest was continuous, covering roughly the same area as today.

The findings of the UKCEH-led modeling study, [published](#) in the journal *npj Biodiversity*, indicate the answer is somewhere in between: woodlands and savannas connected these forest islands. The authors conclude this allowed some animals, which could travel further and were more adaptable, to move between them, while acting as a barrier to more specialist species that rely on a certain type of habitat.

The make-up of an area's vegetation and climate influences [natural evolution](#), meaning that while a species may originally have been present

across a large area, different species gradually evolved in different areas as animals' local environment changed.

Study lead author Dr. Douglas Kelley, a Land Surface Modeler at UKCEH, explains, "Climate changes in the past played a huge role in shaping how the Amazon forest looked and where it grew. Similar changes are coming very soon and much faster.

"Previously, fluctuations in climate took thousands of years to happen, enabling plants and animals to adapt, but now significant global warming is happening within decades due to increasing human-driven greenhouse gas emissions."

The study involved UKCEH, the Ontario Forest Research Institute, Kiel University in Germany, the Met Office in the UK, INPA in Brazil and the Field Museum of Natural History in Chicago.

The team used a combination of advanced climate and vegetation modeling techniques with computer-based predictions of the type of plants that grew during the last Ice Age and their location, based on records of fossilized pollen from sediment. Previous research has largely used single research techniques.

The researchers say, unlike previous studies, they considered the gaps that would have existed between the areas where fossilized pollen has been found and whether animals would have been able to move between habitats. Their advanced computer and statistical modeling indicated what type of vegetation would have grown there, verified by the type of plants recorded in any adjoining areas.

Forest cover was dramatically reduced during the last Ice Age due to the colder, [drier climate](#). There was less carbon dioxide in the atmosphere and the dry conditions restricted tree growth and caused more wildfires

that destroyed many forested areas.

Research co-author Dr. Hiromitsu Sato of the Ontario Forest Research Institute, says, "We believe our study, involving integrated modeling methods, provides the best evidence to date on the origins of the rich biodiversity in the Amazon.

"Using [land surface](#) modeling to provide crucial information on biodiversity is a new and very exciting innovation that could be used in future to investigate history of species where there is a shortage of data."

The researchers' next step is to explore when and where species within groups like monkeys, antbirds, frogs, butterflies, and lianas might have diverged from one another over thousands of years.

**More information:** Douglas I. Kelley et al, Niche-dependent forest and savanna fragmentation in Tropical South America during the Last Glacial Maximum, *npj Biodiversity* (2024). [DOI: 10.1038/s44185-024-00056-4](#)

Provided by UK Centre for Ecology & Hydrology

Citation: Modeling study explains why amazon is such a biodiverse paradise (2024, September 11) retrieved 11 September 2024 from <https://phys.org/news/2024-09-amazon-biodiverse-paradise.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--