


## Organic farms can double plant diversity—but only with time

September 2 2024





Location of the sampled conventional (white circles), young organic (1–11 years since transition, gray circles) and old organic farms (20–35 years since transition, black circles) in southern Sweden. Credit: *Journal of Applied Ecology* (2024). DOI: 10.1111/1365-2664.14650

It takes a long time, up to several decades, before the benefits of organic farming take full effect on land that was previously cultivated conventionally, a new study from Lund University suggests. After thirty years, the plant species richness around the cultivated crop had doubled on organic farms compared to conventional farms.

It is well known that organic farming benefits <u>biodiversity</u> and can mitigate some of the negative effects that arise from farming. What is less known, however, is how long it takes for biodiversity to recover after a transition from conventional to organic farming and how significant the recovery will be.

To seek answers, a research team from Lund University investigated 88 different fields in southern Sweden, on 30 different farms. The results show that after thirty years of organic farming, the number of arable plant species had more than doubled in the fields of the <u>organic farms</u>, from an average of 15 to 35 plant species.

The study is **<u>published</u>** in the *Journal of Applied Ecology*.

"We were surprised by the sharp increase in the number of <u>plant species</u>. After thirty years, we weren't expecting that, and it looks like it may continue to increase," says Romain Carrié, researcher at Lund University who led the study.

The fact that it took so long for the recovery, and that the improvement



in biodiversity continued gradually over several decades, is something that must be kept in mind when evaluating the benefits of organic farming, say the researchers.

"If it takes a while before organic farming's full potential for biodiversity hits, we might have underestimated the benefits that organic methods provide so far. Our results indicate that one must take into account a certain delay after a transition in order not to underestimate the benefit to the environment," says Carrié.

The time aspect must also be included when designing <u>financial support</u> for farmers, he believes, so that they are backed up during the transition phase when yield-increasing ecosystem services, which are dependent on arable plant diversity, are built up.

In the study, it was clear that plants sensitive to herbicides took longer to establish in cereal fields after a switch from conventional agriculture.

"This could mean there is a long-term effect of herbicides even after farmers stop using them on the farm, or that it takes time for these plants to colonize the farm when it has become organic," says Carrié.

At the same time, it was the species that are most sensitive to herbicides that contributed the most to the recovery of the plant community over time in organic farming.

Other factors that affected recolonization were the dispersal ability of the various field plants and their dependence on insect pollination. Greater pollination dependence meant a longer recolonization time. To shorten the <u>recovery time</u> after a conversion to organic farming, it is therefore important to reduce the use of herbicides and to improve conditions for pollinators in conventional agriculture, the researchers summarize in the study.



"We hope that <u>policy makers</u> take note of the conclusions and that we can inspire new policy instruments to stimulate the transition to organic farming in order to preserve biodiversity," says Carrié.

A previous study by other researchers has shown a more immediate recolonization after conversion to <u>organic farming</u>. At that time, however, they studied field edges, which have a different dynamic, and not the fields themselves.

Yet another study has shown a less extensive increase in biodiversity after conversion, but it did not include the same time span after the conversion.

**More information:** Romain Carrié et al, Sensitivity to agricultural inputs and dispersal limitation determine the response of arable plants to time since transition to organic farming, *Journal of Applied Ecology* (2024). DOI: 10.1111/1365-2664.14650

Provided by Lund University

Citation: Organic farms can double plant diversity—but only with time (2024, September 2) retrieved 4 September 2024 from <u>https://phys.org/news/2024-09-farms-diversity.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.