


Study sheds light on potential impacts of bisphenol contaminants on marine appendicularians

September 10 2024



Potential impacts of bisphenol A (BPA) on *Oikopleura dioica* morphology, population fitness, and transcriptomics. Credit: IOCAS

Bisphenol compounds (BPs), crucial ingredients in the production of various polymers, are ubiquitous in our daily lives and industrial

processes. Bisphenol A (BPA), a prevalent BP in the environment, has been extensively studied for its impacts on diverse species, yet its specific mechanisms of action in marine appendicularians remain largely unknown.

Recently, the research team led by Prof. Zhang Guangtao of the Jiaozhou Bay Station at the Institute of Oceanology of the Chinese Academy of Sciences (IOCAS), examined the comprehensive impacts of BPA on the [survival rates](#), growth patterns, and reproductive capabilities of the globally ubiquitous marine appendicularian, *Oikopleura dioica*, at environmentally relevant concentrations.

The study was published in the [Journal of Hazardous Materials](#) on Aug. 16.

Acute toxicity testing revealed a remarkable sensitivity of *Oikopleura dioica* to BPA exposure, with a median lethal concentration (LC50) of merely 0.14 mg/L, 1–2 orders of magnitude lower than that of common aquatic models such as *Daphnia magna* and zebrafish.

Appendicularians are considered the second most abundant marine mesozooplankton and serve as an important shortcut that allows rapid energy transfer from producers to consumers in the marine microbial food web.

"Our work underscores the potential of the appendicularian *Oikopleura dioica* as a valuable sentinel for monitoring bisphenol stress in [marine environments](#)," said Li Shuai, first author of the study from IOCAS.

Further morphological and transcriptomic analyses illuminated the specific toxic effects of BPA on *Oikopleura dioica*. At environmentally relevant concentrations of 125 µg/L, BPA significantly disrupted the growth and reproduction processes of the appendicularian, affecting the

normal secretion of digestive enzymes and phospholipase A2, thereby impairing digestive system function and arachidonic acid metabolism.

Additionally, it downregulated the mRNA expression levels of enzymes associated with carbohydrate and energy metabolism. Notably, even at concentrations generally considered safe for the environment (2.5 µg/L), BPA adversely impacted female fecundity and altered male gene expression patterns.

"Our findings not only deepen the understanding of the adverse effects of bisphenol pollutants in aquatic invertebrates with specific reproductive strategies, but also address the need to reassess environmental safety thresholds of bisphenols," said Prof. Zhang, corresponding author of the paper.

More information: Shuai Li et al, Growth and reproductive toxicity of bisphenol A in *Oikopleura dioica* at environmentally relevant concentrations, *Journal of Hazardous Materials* (2024). [DOI: 10.1016/j.jhazmat.2024.135552](https://doi.org/10.1016/j.jhazmat.2024.135552)

Provided by Chinese Academy of Sciences

Citation: Study sheds light on potential impacts of bisphenol contaminants on marine appendicularians (2024, September 10) retrieved 11 September 2024 from <https://phys.org/news/2024-09-potential-impacts-bisphenol-contaminants-marine.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.