


# Solution to a cosmic mystery—the eccentric orbits of trans-Neptunian objects

September 6 2024



Simulation snapshots of model A1. Credit: *Nature Astronomy* (2024). DOI: 10.1038/s41550-024-02349-x

New evidence suggests that billions of years ago, a star may have passed very close to our solar system. As a result, thousands of smaller celestial bodies in the outer solar system outside Neptune's orbit were deflected into highly inclined trajectories around the sun. It is possible that some of them were captured by the planets Jupiter and Saturn as moons.

These findings come from a team of astrophysicists from Forschungszentrum Jülich and Leiden University in the Netherlands. They were published in two studies in the journals [Nature Astronomy](#) and

*The Astrophysical Journal Letters.*


When we think of our solar system, we usually assume that it ends at the outermost known planet, Neptune. "However, several thousand celestial bodies are known to move beyond the [orbit](#) of Neptune," explains Susanne Pfalzner, astrophysicist at Forschungszentrum Jülich.

It is even suspected that there are tens of thousands of objects with a diameter of more than 100 kilometers. "Surprisingly, many of these so-called trans-Neptunian objects move on eccentric orbits that are inclined relative to the common orbital plane of the planets in the solar system."

Together with her Jülich colleague Amith Govind and Simon Portegies Zwart from Leiden University, Susanne Pfalzner has used more than 3,000 [computer simulations](#) to investigate a possible cause of the unusual orbits: could another star have caused the strange orbits of trans-Neptunian objects?

The three astrophysicists found that a distinctive, close flyby of another star can explain the inclined and eccentric orbits of the known trans-Neptunian celestial bodies. "Even the orbits of very distant objects can be deduced, such as that of the dwarf planet Sedna in the outermost reaches of the solar system, which was discovered in 2003.

"And also objects that move in orbits almost perpendicular to the planetary orbits," says Susanne Pfalzner. Such a flyby can even explain the orbits of 2008 KV42 and 2011 KT19—the two [celestial bodies](#) that move in the opposite direction to the planets.



Saturn's moon Phoebe is a prime example of the unusual properties of irregular moons. Like many others, it orbits Saturn in the opposite direction. Credit: NASA / JPL

"The best match for today's [outer solar system](#) that we found with our simulations is a star that was slightly lighter than our sun—about 0.8 solar masses," explains Pfalzner's colleague Amith Govind. "This star flew past our sun at a distance of around 16.5 billion kilometers. That's about 110 times the distance between Earth and the sun, a little less than four times the distance of the outermost planet Neptune."

However, the scientists' most surprising realization was that the flyby of an alien star billions of years ago could also provide a natural explanation for phenomena closer to home. Susanne Pfalzner and her colleagues found that in their simulations, some trans-Neptunian objects were hurled into our solar system—into the region of the outer giant planets Jupiter, Saturn, Uranus and Neptune.

"Some of these objects could have been captured by the giant planets as moons," says Simon Portegies Zwart from Leiden University. "This would explain why the outer [planets](#) of our solar system have two different types of moons."

In contrast to the regular moons, which orbit close to the planet on circular orbits, the irregular moons orbit the planet at a greater distance on inclined, elongated orbits. Until now, there was no explanation for this phenomenon.

"The beauty of this model lies in its simplicity," says Pfalzner. "It answers several open questions about our solar system with just a single cause."

**More information:** Susanne Pfalzner et al, Trajectory of the stellar flyby that shaped the outer Solar System, *Nature Astronomy* (2024). DOI: [10.1038/s41550-024-02349-x](https://doi.org/10.1038/s41550-024-02349-x)

Susanne Pfalzner et al, Irregular Moons Possibly Injected from the Outer Solar System by a Stellar Flyby, *The Astrophysical Journal Letters* (2024). DOI: [10.3847/2041-8213/ad63a6](https://doi.org/10.3847/2041-8213/ad63a6)

Provided by Forschungszentrum Juelich

Citation: Solution to a cosmic mystery—the eccentric orbits of trans-Neptunian objects (2024, September 6) retrieved 6 September 2024 from <https://phys.org/news/2024-09-solution-cosmic-mystery-eccentric-orbits.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.