

Tropical forests face increased soil carbon loss due to climate change

September 3 2024, by Anne M. Stark



From right, Daniela Cusak, LLNL's Karis McFarlane and Andy Nottingham take soil samples from a rainforest. Credit: Alexandra Hedgpeth

Tropical forests account for more than 50% of the global terrestrial carbon sink, but climate change threatens to alter the carbon balance of these ecosystems.

New research by Lawrence Livermore National Laboratory (LLNL)


scientists and colleagues from Colorado State University and the Smithsonian Tropical Research Institute has found that [warming](#) and drying of tropical forest soils may increase soil carbon vulnerability, by increasing degradation of older carbon. The research [appears](#) in *Nature*.

"These findings imply that both warming and drying, by accelerating the loss of older soil carbon or reducing the incorporation of fresh carbon inputs, will intensify soil carbon losses and negatively impact carbon storage in [tropical forests](#) under climate change," said LLNL scientist Karis McFarlane, lead author of the paper.

Tropical forests exchange more CO₂ with the atmosphere than any other terrestrial biome and store nearly one-third of global soil carbon stocks. Tropical terrestrial ecosystems also have the shortest mean residence time for carbon on Earth, as short as 6-15 years, meaning that any change in carbon inputs or outputs (including CO₂ emitted by soil) could have large and relatively rapid consequences for tropical ecosystem carbon balance and carbon-climate feedbacks.

Climate projections suggest a future that will be both warmer and drier for much of the tropics with increasing drought intensity and dry season length for the Neotropics (a region extending from southern Mexico through Central America and northern South America, including the vast Amazon rainforest).

The research, conducted during climate manipulation experiments in tropical forests in Panama, shows that both whole-profile in situ heating of soil by 4 °C and exclusion of 50% of rainfall increased carbon-14 in the CO₂ released by the soil, increasing the average age of the carbon by the equivalent of ~2–3 years.



Study site locations on the Isthmus of Panama. Credit: *Nature Communications* (2024). DOI: 10.1038/s41467-024-51422-6

Importantly, the mechanisms underlying this shift differed between warming and drying. Warming accelerated decomposition of older carbon as increased CO₂ emissions depleted newer carbon. Drying suppressed decomposition of newer carbon inputs and decreased soil CO₂ emissions, thereby increasing the contributions of older carbon to CO₂ release.

"Field and laboratory experiments suggest that climate warming will stimulate a net loss of global soil carbon to the atmosphere, but how climate warming and drying will interact to influence carbon balance in forests and other ecosystems is less clear," McFarlane said.

Most of the previous work in tropical forests only considered total CO₂ flux rates, which are important for determining the overall carbon balance of tropical forests, but are limited in their ability to uncover mechanisms behind observed change. Those mechanisms can be revealed by carbon-14 values, which indicate the average age of the

carbon sources being metabolized and released as CO₂.

"New" or "young" carbon has been fixed from the atmosphere in the last few years while older "decadal aged" carbon is enriched in carbon-14 relative to the current atmosphere. Even older "century" or "millennial-aged" is depleted in carbon-14 relative to the current atmosphere.

In the current study, the team determined how warming and drying impact the amount and age of carbon released as soil CO₂ in two distinct lowland tropical forest areas in Panama that are subject to experimental soil warming or experimental drying. They measured the carbon-14 and carbon-13 isotopes of soil-respired CO₂.

Using LLNL's Center for Accelerator Mass Spectrometry, McFarlane and her team found that soil warming increased the carbon-14 of respired CO₂ during the wet season, indicative of greater release of "bomb" (circa 1963 from underground nuclear testing) carbon under warmed and wet conditions. Specifically, warming stimulated the decomposition of older soil carbon by increasing overall soil CO₂ release, causing a microbial switch in resource use following the depletion of fresh organic matter to older soil carbon. In contrast, drying reduced total soil CO₂ release, but also increased the carbon-14 of respired CO₂ by limiting the delivery of fresh carbon (from leaf litter or roots) to decomposers.

"This limitation of microbial access to fresh carbon explains the shift toward increased contributions of older carbon in total soil CO₂ emissions with warming and drying," McFarlane said. "Our results suggest that climate change will increase the vulnerability of previously stored [soil](#) carbon in tropical forests by stimulating the decomposition and loss of old [carbon](#)."

More information: Karis J. McFarlane et al, Experimental warming

and drying increase older carbon contributions to soil respiration in lowland tropical forests, *Nature Communications* (2024). [DOI: 10.1038/s41467-024-51422-6](https://doi.org/10.1038/s41467-024-51422-6)

Provided by Lawrence Livermore National Laboratory

Citation: Tropical forests face increased soil carbon loss due to climate change (2024, September 3) retrieved 3 September 2024 from <https://phys.org/news/2024-09-tropical-forests-soil-carbon-loss.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--