

Researchers reassess theories on formation of Earth's atmosphere

September 19 2007



Geochemists at Rensselaer Polytechnic Institute are challenging commonly held ideas about how gases are expelled from the Earth. Their theory, which is described in the Sept. 20 issue of the journal *Nature*, could change the way scientists view the formation of Earth's atmosphere and those of our distant neighbors, Mars and Venus. Their data throw into doubt the timing and mechanism of atmospheric formation on terrestrial planets.

Lead by E. Bruce Watson, Institute Professor of Science at Rensselaer, the team has found strong evidence that argon atoms are tenaciously bound in the minerals of Earth's mantle and move through these minerals at a much slower rate than previously thought. In fact, they found that

even volcanic activity is unlikely to dislodge argon atoms from their resting places within the mantle. This is in direct contrast to widely held theories on how gases moved through early Earth to form our atmosphere and oceans, according to Watson.

Scientists believe that shortly after Earth was formed, it had a glowing surface of molten rock extending down hundreds of miles. As that surface cooled, a rigid crust was produced near the surface and solidified slowly downward to complete the now-solid planet. Some scientists have suggested that Earth lost all of its initial gases, either during the molten stage or as a consequence of a massive collision, and that the catastrophically expelled gases formed our early atmosphere and oceans. Others contend that this early “degassing” was incomplete, and that primordial gases still remain sequestered at great depth to this day. Watson’s new results support this latter theory.

“For the ‘deep-sequestration’ theory to be correct, certain gases would have to avoid escape to the atmosphere in the face of mantle convection and volcanism,” Watson said. “Our data suggest that argon does indeed stay trapped in the mantle even at extremely high temperatures, making it difficult for the Earth to continuously purge itself of argon produced by radioactive decay of potassium.”

Argon and other noble gases are tracer elements for scientists because they are very stable and do not change over time, although certain isotopes accumulate through radioactive decay. Unlike more promiscuous elements such as carbon and oxygen, which are constantly bonding and reacting with other elements, reliable argon and her sister noble gases (helium, neon, krypton, and xenon) remain virtually unchanged through the ages. Its steady personality makes argon an ideal marker for understanding the dynamics of Earth’s interior.

“By measuring the behavior of argon in minerals, we can begin to retrace

the formation of Earth's atmosphere and understand how and if complete degassing has occurred," Watson explained.

Watson's team, which includes Rensselaer postdoctoral researcher Jay B. Thomas and research professor Daniele J. Cherniak, developed reams of data in support of their emerging belief that argon resides stably in crystals and migrates slowly. "We realized from our initial results that these ideas might cause a stir," Watson said. "So we wanted to make sure that we had substantial data supporting our case."

The team heated magnesium silicate minerals found in Earth's mantle, which is the region of Earth sandwiched between the upper crust and the central core, in an argon atmosphere. They used high temperature to simulate the intense heat deep within the Earth to see whether and how fast the argon atoms moved into the minerals. Argon was taken up by the minerals in unexpectedly large quantities, but at a slow rate.

"The results show that argon could stay in the mantle even after being exposed to extreme temperatures," Watson said. "We can no longer assume that a partly melted region of the mantle will be stripped of all argon and, by extension, other noble gases."

But there is some argon in our atmosphere--slightly less than 1 percent. If it didn't shoot through the rocky mantle, how did it get into the atmosphere"

"We proposed that argon's release to the atmosphere is through the weathering of the upper crust and not the melting of the mantle," Watson said. "The oceanic crust is constantly being weathered by ocean water and the continental crust is rich in potassium, which decays to form argon."

And what about the primordial argon that was trapped in the Earth

billions of years ago" "Some of it is probably still down there," Watson said.

Because Mars and Venus have mantle materials similar to those found on Earth, the theory could be key for understanding their atmospheres as well.

Watson and his team have already begun to test their theories on other noble gases, and they foresee similar results. "We may need to start reassessing our basic thinking on how the atmosphere and other large-scale systems were formed," he said.

Source: Rensselaer Polytechnic Institute

Citation: Researchers reassess theories on formation of Earth's atmosphere (2007, September 19) retrieved 18 September 2024 from

<https://phys.org/news/2007-09-reassess-theories-formation-earth-atmosphere.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.