

Does hotter mean healthier?

February 3 2009



Here, the relationship between chile peppers' heat level and plant disease resistance is studied. Credit: Soumaila Sanogo

Phytophthora blight, caused by *Phytophthora capsici*, is a major plant disease that affects many crop species worldwide, including chile peppers in New Mexico. Farmers' observations suggested that *Phytophthora capsici* caused less damage in pepper crops of the hot pepper varieties than low-heat pepper varieties.

A study published in the October 2008 issue of *HortScience* by the research team of Mohammed B. Tahboub (postdoctoral researcher), Soumaila Sanogo (plant pathologist and team leader), Paul W. Bosland (chile pepper breeder), and Leigh Murray (statistician) set out to determine whether or not the severity of Phytophthora blight would be

greater in low-heat than in hot chile peppers.

The most effective means for controlling *Phytophthora* blight are chile pepper cultivars that are genetically resistant to the disease. Some resistant lines have been identified, but currently there are no cultivars that are resistant to the blight in all environments.

Chile pepper fruit become infected during prolonged periods of heavy rain and high humidity in flooded and poorly drained fields. Prior to this study, there had been no systematic assessment of the relationship of chile pepper heat level to chile pepper response to *Phytophthora capsici*. If such a connection could be found, information might have been revealed that would assist breeding programs intended for developing disease-resistant cultivars of pepper.

Based on documented field observations in New Mexico, Arizona, and South Carolina, the researchers hypothesized that peppers that produce high-heat fruits would be more resistant to *Phytophthora* blight than low-heat varieties. The study was conducted by observing infection on both the root and fruit of different varieties of peppers included.

The results of the study concluded, however, that there was no relationship between the heat level of the pepper and the plant's resistance to *Phytophthora* blight. For example, while the disease was slowest to develop on the roots of one variety of jalapeño, it was quickest to develop on the fruit of the same plant.

Conversely, the disease was faster to develop on roots and slower on fruit of all other cultivars. As the root of the plant contains no heat-inducing agents but the fruit does, the slow development on the root and rapid development on the fruit of the jalapeño indicates that heat level is not a factor.

The results of this study indicate that factors other than heat level may be involved in fruit response to *Phytophthora capsici*. Genetic differences and cuticle thickness of the plants and fruits are among other issues that could be relevant, but further study is warranted.

The complete study is available on the ASHS HortScience electronic journal web site: [hortsci.ashspublications.org/c...t/abstract/43/6/1846](http://hortsci.ashspublications.org/content/43/6/1846)

Source: American Society for Horticultural Science

Citation: Does hotter mean healthier? (2009, February 3) retrieved 19 September 2024 from <https://phys.org/news/2009-02-hotter-healthier.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.