


Avoiding the hothouse and the icehouse

11 February 2009



The picture shows the maximum ice distribution on the northern hemisphere during the last ice age. By controlling emissions of fossil fuels we may be able to delay the start of the next ice age for 500,000 years, new research shows. Credit: Dr. Martin Jakobsson, Stockholm Geo Visualization Lab

By controlling emissions of fossil fuels we may be able to greatly delay the start of the next ice age, new research from the Niels Bohr Institute at University of Copenhagen concludes. The results have been published in the scientific magazine, *Geophysical Research Letters*.

From an Earth history perspective, we are living in cold times. The greatest climate challenge mankind has faced has been surviving ice ages that have dominated climate during the past million years. Therefore it is not surprising that back in the relatively cold 1970's prominent scientists like Soviet Union climatologist Mikhail Budyko greeted man-made global warming from CO₂ emissions as a way to keep us out of future ice ages. And there are still those around who feel that continued high fossil fuel emissions are good for this reason. But is the extreme global warming that would result from this a reasonable, and indeed necessary, price to pay to keep ice ages at bay?

In a paper published in the scientific journal *Geophysical Research Letters* 'Long time management of fossil fuels to limit global warming and avoid ice age onsets', Professor Gary Shaffer of the Niels Bohr Institute, University of Copenhagen, and also leader of the research team at the Danish Center for Earth System Science (DCESS), outlines a way to keep the Earth out of both Hot- and Icehouses for a half a million years into the future.

Building up ice sheets

Ice ages start when conditions at high northern latitudes allow winter snowfall to persist over the summer for enough years to accumulate and build ice sheets. Such conditions depend mainly on summer solar radiation there and atmospheric CO₂ concentration. This radiation is modulated on time scales of 20.000, 40.000 and 100.000 years by changes in the Earth's orbit and orientation. Critical summer solar radiation for initiating ice sheet growth can be significantly lower for higher atmospheric CO₂ with its greenhouse warming effect.

Professor Shaffer made long projections over the next 500,000 years with the DCESS Earth System Model to calculate the evolution of atmospheric CO₂ for different fossil fuel emission strategies. He also used results of a coupled climate-ice sheet model for the dependency on atmospheric CO₂ of critical summer solar radiation at high northern latitudes for an ice age onset.

The results show global warming of almost 5 degrees Celsius above present for a "business as usual" scenario whereby all 5000 billion tons of fossil fuel carbon in accessible reserves are burned within the next few centuries. In this scenario the onset of next ice age was postponed to about 170,000 years from now.

Carbon can postpone ice age

However, for a management scenario whereby

fossil fuel use was reduced globally by 20% in 2020 and 60% in 2050 (compared to 1990 levels), maximum global warming was less than one degree Celsius above present. Similar reductions in fossil fuel use have been proposed by various countries like Germany and Great Britain.

In this scenario, combustion pulses of large remaining fossil fuel reserves were then tailored to raise atmospheric CO₂ content high and long enough to parry forcing of ice age onsets by summer radiation minima as long as possible. In this way our present equable interglacial climate was extended for about 500,000 years, three times as long as in the "business as usual" case.

Valuable climate regulation

"It appears to be well established that the strong ice ages the Earth has experienced over the past million years were ushered in by declining levels of atmospheric CO₂. Our present atmospheric CO₂ level of about 385 parts per million is already higher than before the transition to these ice ages" Professor Shaffer notes and adds that "The Earth's orbit is nearly circular at present meaning that the present minimum in summer radiation at high northern latitudes is not very deep. We have already increased atmospheric CO₂ enough to keep us out of the next ice age for at least the next 55,000 years for this orbital setup".

He concludes that "Fossil fuel reserves may be too valuable for regulating climate far into the future to allow the reserves to be consumed within the next few centuries. The price of extreme global warming to avoid ice ages is a high and indeed unnecessary price to pay."

More information: *Geophysical Research Letters*:
www.agu.org/pubs/crossref/2009/2008GL036294.shtml

Source: University of Copenhagen

APA citation: Avoiding the hothouse and the icehouse (2009, February 11) retrieved 18 January 2022 from <https://phys.org/news/2009-02-hothouse-icehouse.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.