

# High construction cost for cycads

July 23 2009

---



A leaf miner insect and an armored scale insect attack the old foliage of *Cycas micronesica*. The damage by these invasive insects does not allow these expensive leaves enough time to pay back the investment into their construction. Credit: Thomas E. Marler

Self-sustaining organisms like plants possess the ability to synthesize their own food using inorganic materials. Plants use water and carbon dioxide to begin this process in their green tissues. The leaf is the organ most often used by plants for this food synthesis. The Western Pacific Tropical Research Center is contributing to the general understanding of

the strategies that plants employ to carry out this foundational step of the food web.

Thomas Marler recently published the results of his research that reveal some of the attributes of the long-lived leaves of Guam's *Cycas micronesica* plants. "These leaves are relatively large and are constructed of tough tissues," said Marler. "Both of these factors indicate the construction costs of this leaf form are substantial."

Green leaves must pay back their own construction costs, but each leaf is also required to supply the energy needed to maintain its live tissues throughout its lifespan. Additionally, large plants like Guam's *Cycas* have an abundance of non-green tissue that requires energy to maintain. Therefore, large plants with this leaf construction strategy must also possess a payback strategy that allows recovery of these costs coupled with an excess in food synthesis to support plant growth.

"These phenomena are understood using a financial analogy where borrowing a modest amount of money for one investment strategy is much easier to pay back and less risky than borrowing a substantial sum of money for a different investment strategy," said Marler. "The manufacture of food by the leaf is analogous to the income generated from the investments."

Results of the research indicate these plants approach the substantial leaf construction investments by delaying the decline in food synthesis capacity that accompanies the leaf aging process. Many fast-growing species, for example, constantly replace their leaves. For these species, food synthesis may be reduced to negligible levels in less than one month after leaf construction. In contrast, the decline in food synthesis for leaves of Guam's *Cycas* is minimal, with rates of two-year-old leaves sustained at 80% of the initial maximum. Marler indicates this sluggish pace of decline is 65-80 times slower than other species for which this

trait has been determined.

The financial risk analogy illuminates a critical environmental issue because Guam's *Cycas* population is being threatened by alien species that feed on its leaves. Guam's alien deer and several alien insects halt a leaf's payback process such that construction costs are not recovered before the leaves are eaten. As the process is repeated the plant effectively starves to death.

This work continues to propel forward the *Cycas* research that is based in Guam. The general lack of physiology research on *Cycas* species is a limitation to our understanding of this important plant group. Every comprehensive written text on *Cycas* makes a point of how little is known about how these plants function. Thomas Marler, through his work at the Western Pacific Tropical Research Center, is busy addressing this void in the literature.

Source: University of Guam

Citation: High construction cost for cycads (2009, July 23) retrieved 21 September 2024 from <https://phys.org/news/2009-07-high-cycads.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.