

Will Kepler find habitable moons?

3 September 2009



Artist's impression of a hypothetical exomoon in orbit around a Saturn-like planet in another planetary system. Image: Dan Durda

(PhysOrg.com) -- Since the launch of the NASA Kepler Mission earlier this year, astronomers have been keenly awaiting the first detection of an Earth-like planet around another star. Now, in an echo of science fiction movies a team of scientists led by Dr David Kipping of University College London thinks that they may even find habitable 'exomoons' too. The new results will appear in a paper in *Monthly Notices of the Royal Astronomical Society*.

Kepler's primary mission is to monitor thousands of stars looking for characteristic dips in their brightness as orbiting planets pass in front of them in so-called 'transit' events. The orbiting observatory should be able to time these transits to an extremely high accuracy.

Dr Kipping has already devised a method for detecting exomoons but no-one was sure whether it could really be used with current technology. He and his team have now modelled the properties of the instruments on Kepler, simulating the expected signal strength that a habitable moon would generate. An exomoon's gravity tugs on the planet it orbits, making the planet wobble during its orbit around its host star. The resulting changes in the position and velocity of the planet should be detectable by Kepler through accurate timing of the

transits.

The scientists considered a wide range of possible planetary systems and found that a fluffy Saturn-like planet (the ringed world is extremely low in mass for its size) gives the best possible chance for detecting a moon, rather than a denser Jupiter-like world. This is because [planets](#) like Saturn are large - blocking out a lot of light as they pass in front of their star - but very light, meaning they will wobble much more than a heavy planet.

If the Saturn-like planet is at the right distance from its star, then the temperature will allow [liquid water](#) to be stable on any sufficiently large moons in orbit around it and these could then be habitable.

The team found that habitable exomoons down to 0.2 times the mass of the Earth are readily detectable with Kepler. Potentially the observatory could look for Earth-mass habitable moons around 25,000 stars up to 500 light-years away from the Sun. In the whole sky, there should be millions of stars which could be surveyed for habitable exomoons with present technology.

Whether or not such bodies are common in the Galaxy is unknown but astronomers now have the tools and the methodology to find out.

Dr Kipping says, "For the first time, we have demonstrated that potentially habitable moons up to hundreds of light years away may be detected with current instrumentation"

'As we ran the simulations, even we were surprised that moons as small as one-fifth of the Earth's mass could be spotted.

'It seems probable that many thousands, possibly millions, of habitable exomoons exist in the Galaxy and now we can start to look for them.'

More information: "On the detectability of habitable moons with Kepler-class photometry", Kipping D. M., Fossey S. J., Campanella G., *Monthly Notices*

of the Royal Astronomical Society, in press. A
preview version can be found at
xxx.lanl.gov/abs/0907.3909

Provided by Royal Astronomical Society ([news](#) :
[web](#))

APA citation: Will Kepler find habitable moons? (2009, September 3) retrieved 27 September 2020 from
<https://phys.org/news/2009-09-kepler-habitable-moons.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.