


Magnetic monopoles detected in a real magnet for the first time


3 September 2009



This is an impression of a "spin spaghetti" of Dirac strings. Credit: HZB / D.J.P. Morris & A. Tennant

Researchers from the Helmholtz Centre Berlin, in cooperation with colleagues from Dresden, St. Andrews, La Plata and Oxford, have for the first time observed magnetic monopoles and how they emerge in a real material. They publish this result in the journal *Science* within the *Science Express* web site on Sept. 3.

Magnetic monopoles are hypothetical particles proposed by physicists that carry a single magnetic pole, either a magnetic North pole or South pole. In the material world this is quite exceptional because [magnetic particles](#) are usually observed as dipoles, north and south combined. However there are several theories that predict the existence of monopoles. Among others, in 1931 the physicist Paul Dirac was led by his calculations to the conclusion that magnetic monopoles can exist at the end of tubes - called Dirac strings - that carry magnetic field. Until now they have remained undetected.



This is a schematic diagram of the neutron scattering experiment: Neutrons are fired towards the sample, and when a magnetic field is applied the Dirac strings align against the field with magnetic monopoles at their ends. The neutrons scatter from the strings providing data which show us the strings properties. Credit: HZB / D.J.P. Morris & A. Tennant

Jonathan Morris, Alan Tennant and colleagues (HZB) undertook a neutron scattering experiment at the Berlin research reactor. The material under investigation was a single crystal of Dysprosium [Titanate](#). This material crystallises in a quite remarkable geometry, the so called pyrochlore-lattice. With the help of neutron scattering Morris and Tennant show that the magnetic moments inside the material had reorganised into so-called „Spin-Spaghetti“. This name comes from the ordering of the dipoles themselves, such that a network of contorted tubes (Strings) develops, through which magnetic flux is transported. These can be made visible by their interaction with the neutrons which themselves carry a [magnetic moment](#). Thus the neutrons scatter as a reciprocal representation of the Strings.

During the neutron scattering measurements a magnetic field was applied to the crystal by the researchers. With this field they could influence the symmetry and orientation of the strings. Thereby it

was possible to reduce the density of the string networks and promote the monopole dissociation. As a result, at temperatures from 0.6 to 2 Kelvin, the strings are visible and have magnetic monopoles at their ends.



Pictured are Bastian Klemke and Jonathan Morris at instrument E2 of the Research-Reactor at HZB in Berlin (Flat-Cone Single Crystal Diffractometer). Credit: HZB / A. Rouvière

The signature of a gas made up by these monopoles has also been observed in heat capacity measured by Bastian Klemke (HZB). Providing further confirmation of the existence of monopoles and showing that they interact in the same way as electric charges.

In this work the researchers, for the first time, attest that monopoles exist as emergent states of matter, i.e. they emerge from special arrangements of dipoles and are completely different from the constituents of the material. However, alongside this fundamental knowledge, Jonathan Morris explains the further meaning of the results: „We are writing about new, fundamental properties of matter. These properties are generally valid for materials with the same topology, that is for magnetic moments on the pyrochlore lattice. For the development of new technologies this can have big implications. Above all it signifies the first time

fractionalisation in three dimensions is observed."

More information:

- Article in *Science Express* 3-Sep-2009: "Dirac Strings and Magnetic Monopoles in Spin Ice Dy₂Ti₂O₇" DOI: 10.1126/science.1178868; D.J.P. Morris, D.A. Tennant, S.A. Grigera, B. Klemke, C. Castelnovo, R. Moessner, C. Czter-nasty, M. Meissner, K.C. Rule, J.-U. Hoffmann, K. Kiefer, S. Gerischer, D. Slobinsky, and R.S. Perry

- Castelnovo, C. et al.: [Magnetic monopoles in spin ice](#). In: *Nature* 451, 42-45, 2008.

Source: Helmholtz Association of German Research Centres ([news](#) : [web](#))

APA citation: Magnetic monopoles detected in a real magnet for the first time (2009, September 3)
retrieved 18 September 2019 from <https://phys.org/news/2009-09-magnetic-monopoles-real-magnet.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.