

First horned dinosaur from Mexico

May 28 2010



This is a fleshed-out artist's rendering of the Mexican horned dinosaur *Coahuilaceratops*. Credit: Lukas Panzarin for the Utah Museum of Natural History

A new species of horned dinosaur unearthed in Mexico has larger horns than any other species - up to 4 feet long - and has given scientists fresh insights into the ancient history of western North America, according to a research team led by paleontologists from the Utah Museum of Natural History at the University of Utah.

"We know very little about the dinosaurs of Mexico, and this find increases immeasurably our knowledge of the dinosaurs living in Mexico during the Late Cretaceous," said Mark Loewen, a paleontologist with the Utah Museum of Natural History and lead author of the study.

The 72-million-year-old rhino-sized creature - *Coahuilaceratops*

magnacuerna - was a four- to five-ton plant-eater belonging to a group called horned dinosaurs, or ceratopsids. The name *Coahuilaceratops magnacuerna* (Koh-WHE-lah-SARA-tops mag-NAH-KWER-na), refers to the Mexican state of Coahuila where it was found, and to the Greek word "ceratops" meaning "horned face." The second part of the name, *magnacuerna*, is a combination of Latin and Spanish meaning "great horn," in reference to the huge horns above the eyes of this dinosaur.

The study, partially funded by the National Geographic Society, was conducted by Mark Loewen, Scott Sampson, Eric Lund and Mike Getty, paleontologists at the Utah Museum of Natural History. Also involved were Andrew Farke of the Raymond M. Alf Museum in Claremont, Calif.; Martha Aguillón-Martínez, Claudio de Leon and Rubén Rodríguez-de la Rosa from the Museum of the Desert in Saltillo, Mexico; and David Eberth of the Royal Tyrrell Museum of Palaeontology in Alberta, Canada.

The new species is to be announced in the book "New Perspectives on Horned Dinosaurs" to be released next week by Indiana University Press.


A Different World

For most of the Late Cretaceous Period, from 97 million to 65 million years ago, high global sea levels resulted in flooding of the central, low-lying portion of North America. As a result, a warm, shallow sea extended from the Gulf of Mexico to the Arctic Ocean, splitting the continent into eastern and western landmasses.

ˆDinosaurs living on the narrow, peninsula-like western landmass - known as Laramidia - occupied only a narrow belt of plains that were sandwiched between the seaway to the east and rising mountains to the west. Central America had not formed at the time, which made Mexico the southern tip of this island continent.

In many ways, the Late Cretaceous is the best-understood time during the Age of Dinosaurs, thanks in large part to more than 120 years of dinosaur hunting in Canada, Montana, New Mexico and the Dakotas. Recent work has revealed new dinosaurs living at the same time in Utah, New Mexico and Texas, yet the dinosaurs from Mexico have remained virtually unknown.

"As the southernmost dinosaurs on Laramidia, we are confident that Mexican dinosaurs will be a critical element in unraveling the ancient mystery of this island continent," Sampson said.



Fossils of a new species of horned dinosaur -- 72-million-year-old *Coahuilaceratops magnacuerna* -- excavated in southern Mexico are shown superimposed on a line drawing of the creature's skull. Credit: Utah Museum of Natural History, University of Utah

Loewen described the arid, desert terrain where the dinosaur was recovered as nothing like Mexico during the Late Cretaceous. About 72 million years ago, the region was a humid estuary with lush vegetation, an area where salt water from the ocean mixed with fresh water from rivers, much like the modern Gulf Coast of the southeastern United

States. Many dinosaur bones in the area are covered with fossilized snails and marine clams, indicating that the dinosaurs inhabited environments adjacent to the seashore.

The rocks in which *Coahuilaceratops* was found also contain large fossil deposits of jumbled duck-bill dinosaur skeletons. These sites appear to represent mass death events, perhaps associated with storms such as hurricanes that occur in the region today.

"Sitting near the southern tip of Laramidia, this region may have been hammered by monstrous storms," Sampson said. "If so, such periodic cataclysms likely devastated miles of coastline, killing off large numbers of dinosaurs."

Recovering a Giant Horned Head

Until recent years, there have been few large-scale paleontological projects in Mexico focused on the Mesozoic Era, from 253 million to 65 million years ago, also known as the Age of Dinosaurs. Indeed *Coahuilaceratops* is among the first dinosaurs from Mexico to be named.

Sampson spearheaded the paleontological expeditions to Coahuila in 2002 and 2003, securing funds from the University of Utah and National Geographic Society.

Coahuilaceratops comes from a rock unit known as the Cerro del Pueblo Formation, which dates to between 71.5 million and 72.5 million years ago. The skeletons, which de Leon discovered in 2001 near the town of Porvenir de Jalpa, approximately 40 miles west of Saltillo, were excavated in 2003. The fossils then were prepared at the Utah Museum of Natural History, requiring two years of meticulous work by skilled volunteer preparator Jerry Golden.

Based on the bone development of the skull and skeleton, the scientists believe that this animal was an adult at the time of death. Remains of a juvenile animal of the same species were also found at the site.

Coahuilaceratops was about 22 feet long as an adult, 6 feet to 7 feet tall at the shoulder and hips, with a 6-foot-long skull, and likely weighed about four to five tons.

"Being one of the largest herbivores in its ecosystem, adult *Coahuilaceratops* probably didn't have to worry about large tyrannosaur predators," Farke said.

By far the most obvious characteristic of *Coahuilaceratops* is its massive pair of horns, one above each eye. While the researchers lack a complete horn, they estimate from fossils they excavated that the horns were 3 feet to 4 feet long, Loewen said.

Although such horns are common features of ceratopsid dinosaurs, those of *Coahuilaceratops* appear to be the largest known for the group, exceeding the size of eye horns even in Triceratops. Scientists are uncertain of the massive eye horns' purpose, but the most widely accepted idea is that they were related to reproductive success, functioning to attract mates and fight with rivals of the same species.

Loewen explained that *Coahuilaceratops* represents the first occurrence of an identifiable species of horned dinosaur in southern Mexico. "The horned dinosaurs are an extraordinary example of vertebrate evolution," he said. They evolved and diversified on Laramidia along a thin strip of land that stretched from Alaska to Mexico. "Finding this horned dinosaur so far south in Mexico offers us a different picture of what the ancestors of Triceratops were like."



This is a line drawing of the skull of a new species of horned dinosaur unearthed in southern Mexico. Credit: Utah Museum of Natural History, University of Utah

An Ancient Ecosystem Revealed

In addition to *Coahuilaceratops*, the research team found remains of two other horned dinosaurs, which are less well understood. "We need more material to figure out what these other horned dinosaurs looked like," Getty said.

The latest expedition also recovered remains of two duck-bill dinosaurs, as well as the remains of carnivores, including large tyrannosaurs (smaller, older relatives of T. rex) and more diminutive Velociraptor-like predators armed with sickle-claws on their feet.

Together with an abundance of fossilized bones, researchers discovered the largest assemblage of dinosaur trackways known from Mexico, an extensive area crisscrossed with the tracks of different kinds of dinosaurs. In all, the emerging picture shows a diverse group of dinosaurian herbivores and carnivores, perhaps representing a previously unknown assemblage of species.

"Rather than focusing only on individual varieties of dinosaurs, we are attempting to reveal what life was like in Mexico 72 million years ago, and understand how the unique ecosystem of Mexico relates to ecosystems to the north at the time," said Loewen.

Few North American dinosaurs from this time period are known outside of the Drumheller region of Alberta. Eberth explained that researchers now have two points of comparison to examine not only different dinosaurs, but also different environments and ecologies.

As might be suspected, [paleontologists](#) are excited about the future paleontological potential of this area. "There are definitely more dinosaurs to be discovered in the region," Lund said.

"[Dinosaurs](#) from this particular period are important because this is a time that is relatively poorly understood," said Don Brinkman, a researcher at the Royal Tyrrell Museum. He is studying non-dinosaur vertebrates found at the site, including turtles, fish, and lizards. "The locality in [Mexico](#) goes a long way to filling in a gap in our knowledge of the record of changes in dinosaur assemblages throughout the Late Cretaceous."

Provided by University of Utah

Citation: First horned dinosaur from Mexico (2010, May 28) retrieved 24 April 2024 from <https://phys.org/news/2010-05-horned-dinosaur-mexico.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.